

11-09-00

A

PATENT  
Attorney Docket No.: MSU-04769

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

## Box Patent Application


Assistant Commissioner For Patents  
Washington, D.C. 20231

## NEW APPLICATION TRANSMITTAL

Transmitted herewith for filing is the patent application of Christoph Benning, Sherrie Lea Sanda and Bin Yu for Compositions And Methods For The Synthesis And Subsequent Modification Of Uridine-5'-Diphosphosulfoquinovose (UDP-SQ).

## CERTIFICATION UNDER 37 C.F.R. § 1.10

I hereby certify that this New Application Transmittal and the documents referred to as enclosed therein are being deposited with the U.S. Postal Service on this date **November 8, 2000** in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number **EL512052511US** addressed to: **Box Patent Application**, Assistant Commissioner For Patents, Washington, D.C. 20231.



Traci E. Light

## 1. Type Of Application

This new application is for a(n)

☒ Original (nonprovisional)

## 2. Papers Enclosed That Are Required For Filing Date Under 37 C.F.R. § 1.53(b) (Regular) or 37 C.F.R. § 1.153

## (Design) Application

79 Pages of Specification

3 Pages of Claims

1 Page of Abstract

18 Sheets of Informal Drawings

## 3. Declaration

☒ Enclosed☒ Unexecuted.

## 4. Inventorship Statement

The inventorship for all the claims in this application is:

☒ the same

## 5. Language

☒ English

## 6. Assignment

☒ An assignment of the invention to **Michigan State University** is attached.☒ Form PTO-1595 will follow.

## 7. Fee Calculation (37 C.F.R. § 1.16)

☒ Regular application

## CLAIMS AS FILED

Number Filed	Number Extra	Rate	Basic Fee - \$690.00 (37 C.F.R. § 1.16(a))
Total Claims (37 C.F.R. § 1.16(c))	14 - 20 =	0 × \$18.00 =	\$0.00
Independent Claims (37 C.F.R. § 1.16(b))	3 - 3 =	0 × \$78.00 =	\$0.00
Multiple Dependent Claim(s), if any (37 C.F.R. § 1.16(d))	+ \$260.00 =		\$0.00
Filing Fee Calculation			\$690.00

## 8. Small Entity Statement(s)

☒ Verified Statement(s) that this is a filing by a small entity under 37 C.F.R. §§ 1.9 and 1.27 is(are) attached.

Filing Fee Calculation (50% of above) \$345.00

9. **Fee Payment Being Made At This Time**

☒ Enclosed  
☒ basic filing fee \$345.00  
**Total Fees Enclosed \$345.00**

10. **Method of Payment of Fees**

☒ Check in the amount of \$345.00

11. **Authorization To Charge Additional Fees and Credit Overpayment**

☒ The Commissioner is hereby authorized to charge payment of any fees associated with this communication or credit any overpayment to Deposit Account No.: **08-1290**. An originally executed duplicate of this transmittal is enclosed for this purpose.


12. **Power of Attorney by Assignee**

☒ Enclosed

13. **Return Receipt Postcard**

☒ Enclosed

Dated: November 8, 2000

  
\_\_\_\_\_  
Peter G. Carroll  
Registration No.: 32,837

MEDLEN & CARROLL, LLP  
220 Montgomery Street, Suite 2200  
San Francisco, California 94104  
(415) 705-8410

☒ **Statement Where No Further Pages Added**

☒ This transmittal ends with this page.

Applicant / Patentee: Christoph Benning *et al.*  
For: **Compositions And Methods For The Synthesis And Subsequent  
Modification Of Uridine-5'-Diphosphosulfoquinovose (UDP-SQ)**

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS  
(37 CFR §§ 1.9(f) and 1.27(d) - NONPROFIT ORGANIZATION)**

I hereby declare that I am an official empowered to act on behalf of the university (or other institution of higher education) identified below:

Michigan State University  
238 Administration Building, East Lansing, Michigan 48824-1046

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR § 1.9(e) for purposes of paying reduced fees under §§ 41(a) and (b) of Title 35, United States Code with regard to the invention entitled **Compositions And Methods For The Synthesis And Subsequent Modification Of Uridine-5'-Diphosphosulfoquinovose (UDP-SQ)**, by inventors named **Christoph Benning *et al.***, described in the specification filed herewith.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR § 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Date: \_\_\_\_\_

By: \_\_\_\_\_

Name: \_\_\_\_\_

Title: \_\_\_\_\_

09709020 410800

**COMPOSITIONS AND METHODS FOR THE SYNTHESIS AND  
SUBSEQUENT MODIFICATION OF URIDINE-5'-  
DIPHOSPHOSULFOQUINOVOSE (UDP-SQ)**

**FIELD OF THE INVENTION**

5           The present invention relates to compositions and methods for the synthesis and subsequent modification of uridine-5'-diphosphosulfoquinovose (UDP-SQ). The methods of the present invention comprise the utilization of recombinant enzymes from *Arabidopsis thaliana*, UDP-glucose, and a sulfur donor to synthesize UDP-SQ, and the subsequent modification of UDP-SQ to form compounds including, but not  
10       limited to, 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG) and alkyl sulfoquinovoside.

**BACKGROUND**

          Uridine-5'-diphosphosulfoquinovose (UDP-SQ) is a unique sugar nucleotide which carries a negative charge at its sulfonate group. UDP-SQ is believed to react with sugar nucleotide-dependent glycosyltransferases and donate its sulfonate group to  
15       other substrates in order to form valuable compounds including, but not limited to, 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG). UDP-SQ is thought to be the direct precursor of SQDG, to which it donates its unique sulfonic acid head group, sulfoquinovose. However, there is not a simple, rapid method of synthesizing UDP-SQ, or an efficient method for subsequent modification of UDP-SQ to compounds  
20       including, but not limited to, SQDG and alkyl sulfoquinovoside.

          SQDG is an abundant sulfur-containing non-phosphorous glycerolipid that is specifically associated with photosynthetic (thylakoid) membranes of higher plants, mosses, ferns, algae, and most photosynthetic bacteria. SQDG is universally associated with oxygenic photosynthesis and is an important component of the  
25       biological sulfur cycle.

          SQDG has also been shown to be a potent inhibitor of several mammalian DNA polymerases and Human Immunodeficiency Virus Reverse Transcriptase 1 (HIV-RT1), and as such, is valuable as an anti-viral compound. (Ohta *et al.*,



"Sulfoquinovosyldiacylglycerol, KM043, a new potent inhibitor of eukaryotic DNA polymerases and HIV-reverse transcriptase type 1 from a marine red alga, *Gigartina tenella*," *Chem. Pharm. Bull.*, 46(4): 684-86 (1998)). Moreover, SQDG has also been demonstrated to be valuable due to its anti-tumor promoting properties and its ability to enhance the cytotoxic effects of anti-cancer chemotherapy agents. (Shirahashi *et al.*, "Isolation and Identification of Anti-tumor-Promoting Principles from the Fresh-Water Cyanobacterium *Phormidium tenue*," *Chem. Pharm. Bull.*, 41(9): 1664-66 (1993)). Furthermore, SQDG is commonly thought to have excellent detergent properties. (Benson, A.A., "The Plant Sulfolipid," *Adv. Lipid Res.*, 1: 387-94 (1963)). Thus, a method of producing UDP-SQ, and its subsequent modification to compounds including, but not limited to, SQDG, is desirable.

Traditionally, UDP-SQ has been synthesized through a series of chemical reactions. (Heinz *et al.*, "Synthesis of different nucleoside 5'-diphospho-sulfoquinovoses and their use for studies on sulfolipid biosynthesis in chloroplasts," *Eur. J. Biochem.*, 184: 445-453 (1989)). However, this chemical production is highly involved, results in low yields of UDP-SQ, and requires several days to complete. (*Id.*) Moreover, previous studies of SQDG required time-consuming isolation and purification of the anionic sulfolipid from photosynthetic organisms. (Ohta *et al.*, "Action of a New Mammalian DNA Polymerase Inhibitor, Sulfoquinovosyl diacylglycerol," *Biol. Pharm. Bull.*, 22(2): 111-16 (1999); Gustafson *et al.*, "AIDS-Antiviral Sulfolipids From Cyanobacteria (Blue-Green Algae)," *J. Natl. Cancer Inst.*, 81: 1254-258 (1989)). Thus, what is needed is a more simple, rapid method of synthesizing UDP-SQ, and for the subsequent modification of UDP-SQ to compounds including, but not limited to, SQDG.

## SUMMARY OF THE INVENTION

The present invention relates to methods for the synthesis and subsequent modification of uridine-5'-diphosphosulfoquinovose (UDP-SQ). The methods of the present invention comprise the utilization of recombinant enzymes from *Arabidopsis*

*thaliana*, UDP-glucose, and a sulfur donor to synthesize UDP-SQ. Unlike the current methods for the synthesis of UDP-SQ, the synthesis methods of the present invention are simple and rapid. Indeed, the production of UDP-SQ by the methods of the present invention can be completed in less than an hour.

5 In one embodiment, the present invention contemplates a method for synthesizing UDP-SQ comprising: a) providing: i) uridine-5'-diphosphoglucose (UDP-Glc); ii) a sulfur donor; and iii) a peptide capable of catalyzing the conversion of UDP-Glc to uridine-5'-diphosphosulfoquinovose (UDP-SQ); and b) reacting said UDP-Glc with said first peptide and said sulfur donor under such conditions that UDP-SQ is  
10 generated.

It is not intended that the present invention be limited by any specific first peptide capable of catalyzing the conversion of UDP-Glc and a sulfur donor to UDP-SQ. In one embodiment, said first peptide is SQD1, a gene product encoded by the nucleic acid sequence set forth in SEQ ID NO: 6.

15 It is not intended that the present invention be limited by the use of any specific sulfur donor. In one embodiment, the sulfur donor is selected from a group comprising sulfate, sulfite, sulfide, thiosulfate, sulfoglutathione, adenosine 5'-phosphosulfate (APS), and 3'-phosphoadenosine-5'-phosphosulfate (PAPS). In a preferred embodiment, the sulfur donor is sulfite.

20 It is not intended that the present invention be limited by the use of any specific method to express or produce a peptide capable of catalyzing the conversion of UDP-Glc and a sulfur donor to UDP-SQ. In one embodiment, the present invention contemplates the cloning of the *sqd1* gene cDNA into the group of protein expression vectors such as pQE-9, pQE-16, pQE-31, pQE-32, pQE-40, pQE-60, pQE-70, pQE-80,  
25 pQE-81, pQE-82, or pQE-100. In another embodiment, the present invention contemplates the cloning of the *sqd1* gene cDNA into the protein expression vector, pQE-30. (See Figure 3).

The methods of the present invention are conveniently carried out in a reaction vessel or container. It is not intended that the present invention be limited to any

particular reaction vessel. A variety of containers can be used, including but not limited to tubes, flasks and other glassware.

In an alternative embodiment, the invention contemplates the transformation of plant cells or tissues such that the *sqd1* gene product is expressed. In one  
5 embodiment, the present invention contemplates the cloning of the *sqd1* gene cDNA (SEQ ID NO: 6) into a binary vector for introduction into *Agrobacterium tumefaciens*, and the subsequent generation of transgenic plant cells via Agrobacterial transformation.

It is not intended that the present invention be limited by the use of any  
10 specific method to purify a recombinant peptide capable of catalyzing the conversion of UDP-Glc to UDP-SQ. In one embodiment, the present invention contemplates purification of the peptide by use of 6 His-tag incorporated into the protein expression vector that allows protein affinity purification over a nickel-nitriloacetic acid (Ni-NTA) agarose resin-based chromatography column.

It is not intended that the present invention be limited by the use of any  
15 specific method for the detection of UDP-SQ synthesis. The present invention contemplates a variety of method, or assay, formats. In one embodiment, an enzyme assay is provided to measure the conversion of UDP-glucose to UDP-SQ as a reflection of the activity of SQD1. In another embodiment, a coupled adenosine 5'-  
20 phosphosulfate (APS)/SQD1 assay is contemplated.

The present invention relates to methods for the subsequent modification of uridine-5-diphosphosulfoquinovose (UDP-SQ) to synthesize compounds including, but not limited to, 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG). Unlike the current methods for the synthesis of UDP-SQ, the synthesis methods of the present invention  
25 are rapid and simple.

In one embodiment, the present invention contemplates a method for synthesizing UDP-SQ comprising: a) providing: i) uridine-5'-diphosphoglucose (UDP-Glc); ii) a sulfur donor; iii) a first peptide capable of catalyzing the conversion of UDP-Glc to uridine-5'-diphosphosulfoquinovose (UDP-SQ); and iv) a second peptide  
30 capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol; b) reacting

said UDP-Glc with said first peptide and said sulfur donor under such conditions that UDP-SQ is generated; and c) treating said UDP-SQ with said second peptide under conditions such that sulfoquinovose diacylglycerol is generated.

It is not intended that the present invention be limited by the use of any specific second peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol. In one embodiment, said second peptide is a gene product of the nucleic acid sequence set forth in SEQ ID NO: 1 derived from a Cyanobacteria species. In another embodiment, said second peptide is a gene product derived from *Arabidopsis thaliana* and encoded by a nucleic acid sequence selected from the group SEQ ID NO: 3, SEQ ID NO: 4, and SEQ ID NO: 5.

It is not intended that the present invention be limited by the use of any specific method to express or produce a peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol. In one embodiment, the present invention contemplates the cloning of the *sqdX* gene into the group of protein expression vectors comprising pQE-9, pQE-16, pQE-31, pQE-32, pQE-40, pQE-60, pQE-70, pQE-80, pQE-81, pQE-82, pQE-100. In another embodiment, the present invention contemplates the cloning of the *sqdX* gene into the protein expression vector, pQE-30. (See Figure 3). In a further embodiment, the *sqdX* gene is cloned into the protein expression vector pACYC184.

In an alternative embodiment, the invention contemplates the transformation of plant cells or tissues such that the *sqdX* gene product is expressed. In one embodiment, the present invention contemplates the cloning of the *sqdX* gene cDNA (SEQ ID NO: 1) into a binary vector for introduction into *Agrobacterium tumefaciens*, and the subsequent generation of transgenic plant cells via Agrobacterial transformation. In another embodiment, said gene product is encoded by a nucleic acid sequence selected from the group SEQ ID NO: 3, SEQ ID NO: 4, and SEQ ID NO: 5.

It is not intended that the present invention be limited by the use of any specific method to purify a recombinant peptide capable of transferring sulfoquinovose

from UDP-SQ onto diacylglycerol. In one embodiment, the present invention contemplates purification of the peptide by use of 6 His-tag incorporated into the protein expression vector that allows protein affinity purification over a nickel-nitriloacetic acid (Ni-NTA) agarose resin-based chromatography column.

5 It is not intended that the present invention be limited by the use of any specific method for the detection of SQDG synthesis. The present invention contemplates a variety of assay formats. In one embodiment, the synthesis of SQDG is visualized with iodine vapor and identified by co-chromatography with an *Arabidopsis thaliana* leaf lipid extract known to contain SQDG. In another  
10 embodiment, production of SQDG is verified by quantitative analysis wherein reaction products are isolated from the TLC plates and used to prepare fatty acid methyl esters. The methyl esters are quantified by gas chromatography using myristic acid as the internal standard.

15 It is not intended that the invention be limited to the independent expression of a peptide capable of catalyzing the conversion of UDP-Glc and a sulfur donor to UDP-SQ in a single host organism or plant. Moreover, it is also not intended that the invention be limited to the independent expression of a second peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol in a single host organism or plant. In one embodiment, the invention contemplates the co-expression  
20 of both of the peptides described above in a single host organism. In an alternative embodiment, the invention contemplates the transformation of plant cells or tissues such that both peptides are co-expressed.

25 The present invention contemplates a method for the modification of UDP-SQ comprising: a) providing: i) uridine-5'-diphosphoglucose; ii) a sulfur donor; iii) a peptide capable of catalyzing the conversion of uridine-5'-diphosphoglucose to uridine-5'-diphosphosulfoquinovose; iv) an acid catalyst; v) a short-chain alcohol; and vi) a long-chain alcohol; b) reacting said uridine-5'-diphosphoglucose with said peptide and said sulfur donor under such conditions that uridine-5'-diphosphosulfoquinovose is generated; c) reacting said uridine-5'-diphosphosulfoquinovose with said short-chain  
30 alcohol and said acid catalyst under such conditions that a short-chain alkyl

sulfoquinovoside is generated; and d) treating said short-chain alkyl sulfoquinovoside with said long-chain alcohol under such conditions that a long-chain alkyl sulfoquinovoside is generated.

In contrast to current methods for the production of alkyl sulfoquinovoside-like compounds, the method of the present invention produces a group of substances consisting of a glycosidic unit sulfonated at the C-6 position and acetalized at the C-1 position with an alcohol. Moreover, the alkyl sulfoquinovosides produced by the present invention, unlike other anionic surface-active agents, can be obtained from renewable natural resources and are biodegradable.

It is not intended that the invention be limited by the short-chain alcohol chosen for the method. In one embodiment, the short-chain alcohol is selected from the group methanol, ethanol, propanol, pentanol, hexanol, heptanol, octanol, nonanol, including isomers thereof. In another embodiment, the short-chain alcohol is butanol.

It is not intended that the invention be limited by the acid catalyst chosen for the method. In one embodiment, the acid catalyst is selected from the group comprising  $\text{H}_2\text{SO}_4$ ,  $\text{HCl}$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{BF}_3$ , ortho-toluenesulfonic acid, meta-toluenesulfonic acid, alkylbenzenesulfonic acid, secondary alkyl-sulfonic acid, sulfonic resin, alkylsulfate, alkylbenzenesulfonate, alkyl-sulfonate, and sulfosuccinic acid. In another embodiment, the acid catalyst is para-toluenesulfonic acid.

It is not intended that the invention be limited by the long-chain alcohol chosen for the method. In one embodiment, the long-chain alcohol is a fatty alcohol selected from the group of n-dodecyl alcohol, n-tetradecyl alcohol, n-octadecyl alcohol, n-octyl alcohol, n-decyl alcohol, undecyl alcohol, and tridecyl alcohol. In another embodiment, the long-chain alcohol is a technical mixture of about 3 parts by weight lauryl alcohol and 1 part by weight myristyl alcohol. In another embodiment, the long-chain alcohol is a branched-chain primary alcohol including, but not limited to, oxaalcohol. In another embodiment, the long-chain alcohol is n-hexadecyl alcohol.

It is not intended that the invention be limited by the alkyl sulfoquinovoside produced by the method. The present invention contemplates the production of a variety of alkyl sulfoquinovosides and mixtures thereof. In one embodiment,

the alkyl sulfoquinovosides produced are comprised of a mixture of short and long-chain alkyl sulfoquinovosides. In another embodiment, alkyl oligosulfoquinovosides are produced. In another embodiment, alkyl polysulfoquinovosides are produced. In a further embodiment, alkyl monosulfoquinovosides are produced.

5           The present invention also relates to compositions utilized in the biosynthesis of UDP-SQ and its subsequent modification to compounds including, but not limited to, SQDG and alkyl sulfoquinovoside. In one embodiment, the composition is a substantially pure nucleotide sequence comprising at least a portion of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5. In another embodiment, the  
10       composition comprises RNA transcribed from at least a portion of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5. In another embodiment, the composition comprises protein translated from the RNA transcribed from at least a portion of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5. In  
15       another embodiment, the composition comprises antibodies produced from the translated protein. In a further embodiment, the composition comprises expression constructs comprising at least a portion of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5. In another embodiment, the composition comprises transgenic plant cells or tissues comprising at least a portion of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5.

## 20       **DESCRIPTION OF THE DRAWINGS**

Figure 1 schematically shows the biochemical pathway for UDP-SQ biosynthesis.

Figure 2 is a chromatograph showing the results of an assay to detect the conversion of UDP-Glc by SQD1. The chromatographic analysis of <sup>14</sup>C-labeled  
25       substrate and reaction products by HPLC is shown (A-C). (A) UDP-Glc without SQD1 protein, (B) UDP-Glc and SQD1 protein, (C) UDP-Glc, SQD1 protein, and sulfite, (D) authentic <sup>35</sup>S-labeled UDP-SQ isolated from the *sqdD* mutant of the cyanobacterium, *R. sphaeroides*. U1 and U2, products as described in the text.

Figure 3 schematically shows the vector maps, including restriction endonuclease recognition sites, of the protein expression vectors pQE-30, pQE-31, and pQE-32.

Figure 4 is a photograph of a sodiumdodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) gel showing results of the purification of recombinant SQD1. SDS-PAGE analysis of (A) crude *E. coli* cell culture extract expressing SQD1 protein and Ni-NTA column purification of (B) SQD1 and (C) Thr145Ala mutant. (4 µg of each).

Figure 5 schematically shows the biochemical pathway for SQDG biosynthesis involving the transfer of sulfoquinovose onto Diacylglycerol (DAG).

Figure 6 schematically shows the vector map, including restriction endonuclease recognition sites, of the protein expression vector pACYC184. This plasmid is a small, low copy-number *E. coli* cloning vector that is 4,244 base pairs in length and carries tetracyclin (base numbers 1580-2770) and chloramphenicol-resistance (base numbers 219-3804) genes. The map shows the location of sites for restriction enzymes that cleave the molecule once or twice; unique sites are shown in bold type. The coordinates refer to the position 5' base in each recognition sequence. Nucleotide number 1 of the vector is the first "G" of the unique *EcoRI* site, "GAATTC." The map also shows the relative positions of the antibiotic resistance genes and the origin of DNA replication (ORI) at base numbers 845-847.

Figure 7 schematically shows one embodiment for the chemical modification of UDP-SQ with short and long-chain alcohols and an acid catalyst to produce alkyl sulfoquinovoside.

Figure 8 is a chromatograph showing the results of a coupled APS reductase/SQD1 assay. HPLC chromatograms of reaction products and standards are shown. (A) <sup>35</sup>S-labeled substrate APS without enzymes; (B) <sup>35</sup>S-labeled reaction products following the incubation with APS reductase alone, or (C) in the presence of APS reductase and SQD1; (D) <sup>14</sup>C-labeled UDP-SQ (U2) from the standard SQD1 assay.



Figure 9 shows TLC results of an assay of sulfolipid synthase associated with thylakoid membranes which specifically converts UDP-SQ and diacylglycerol to SQDG. (A) Thin-layer chromatography of lipids following the incubation of spinach thylakoid membranes with labeled reaction product U2 or, for control purposes, <sup>14</sup>C-labeled UDP-Gal the substrate for galactolipid biosynthesis. Lipids were visualized by autoradiography. (B) Iodine staining of the U2 lane. DGDG (digalactosyldiacylglycerol); MGDG (monogalactosyldiacylglycerol); PC, (phosphatidylcholine); PG (phosphatidylglycerol); SQDG (sulfoquinovosyldiacylglycerol).

Figure 10 shows the nucleic acid sequence of the Cyanobacterial *sqdX* gene (SEQ ID NO: 1) (submitted to GenBank data base and assigned accession number U45308, nucleotide numbers 1800-2933). The start and stop codons are highlighted for emphasis.

Figure 11 shows the genomic nucleic acid sequence of *Arabidopsis thaliana* containing the *AtSQDX-1* gene (SEQ ID NO: 3 ) (submitted to GenBank data base and assigned accession number AL137189, nucleotide numbers 82324-85302).

Figure 12 shows the genomic nucleic acid sequence of *Arabidopsis thaliana* containing the *AtSQDX-2* gene (SEQ ID NO: 4) (submitted to GenBank data base and assigned accession number AL021768, nucleotide numbers 1691-4227).

Figure 13 shows the genomic nucleic acid sequence of *Arabidopsis thaliana* containing the *AtSQDX-3* gene (SEQ ID NO: 5) (submitted to GenBank data base and assigned accession number AC008016, nucleotide numbers 114774-117142).

Figure 14 shows the nucleic acid sequence of the *Arabidopsis thaliana SQD1* gene cDNA (SEQ ID NO: 6) (submitted to GenBank data base and assigned accession number AF022082). The start and stop codons are highlighted for emphasis.

Figure 15 shows the nucleic acid sequence of the Cyanobacterial *sqdB* gene (SEQ ID NO: 8) (submitted to GenBank data base and assigned accession number U45308, nucleotide numbers 576-1784). The start and stop codons are highlighted for emphasis.

Figure 16 shows the amino acid sequence of the *Arabidopsis thaliana* *SQD1* gene cDNA product (SEQ ID NO: 7) (submitted to GenBank data base and assigned accession number AF022082).

Figure 17 shows the amino acid sequence of the Cyanobacterial *sqdX* gene product (SEQ ID NO: 2) (submitted to GenBank data base and assigned accession number U45308).

Figure 18 shows the amino acid sequence of the Cyanobacterial *sqdB* gene product (SEQ ID NO: 9) (submitted to GenBank data base and assigned accession number U45308).

## DEFINITIONS

To facilitate understanding of the invention, a number of terms are defined below.

"Associated peptide" as used herein refers to peptides that are bound directly or indirectly to other peptides. Associated peptides that are bound indirectly may have one or more other peptides bound between the two associated peptides. Peptides may be bound via peptide bonds, covalent bonds and non-covalent bonds.

"In operable combination," "in operable order" and "operably linked" as used herein refer to the linkage of nucleic acid sequences in such a manner that a nucleic acid molecule capable of directing the transcription of a given gene and/or the synthesis of a desired protein molecule is produced. The term also refers to the linkage of amino acid sequences in such a manner so that a functional protein is produced.

"Expression construct," "expression vector" and "plasmid" as used herein, refer to one or more recombinant DNA or RNA sequences containing a desired coding sequence operably linked to sequences necessary for the expression of the coding sequence in a cell or host organism (*e.g.*, mammal). The sequence may be single or double stranded.

"Reporter construct," "reporter gene" and "reporter protein" as used herein, refer to DNA or amino acid sequences, as appropriate, that, when expressed in a host cell or organism, may be detected, measured or quantitated.

As used herein, the term "purified" or "to purify" refers to the removal of one or more (undesired) components from a sample. For example, where recombinant polypeptides are expressed in bacterial host cells, the polypeptides are purified by the removal of host cell proteins thereby increasing the percent of recombinant polypeptides in the sample.

As used herein, the term "partially purified" refers to the removal of contaminants of a sample to the extent that the substance of interest is recognizable by techniques known to those skilled in the art (*e.g.*, by staining, blotting, etc.) as accounting for a measurable amount (*e.g.*, picograms, nanograms, micrograms, etc.) in the mixture.

As used herein, the term "substantially purified" refers to molecules, (*e.g.*, nucleic or amino acid sequences) that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free and more preferably 90% free from other components with which they are naturally associated.

As used herein, when a solution passes through the solid support matrix, it comprises the "flow through." Material that does not bind, if present, passes with the solution through the matrix into the flow through. To eliminate all non-specific binding, the matrix is "washed" with one or more wash solutions which, after passing through the matrix, comprise one or more "effluents." "Eluent" is a chemical solution capable of dissociating material bound to the matrix (if any); this dissociated material passes through the matrix and comprises an "eluate."

"Antibody" as used herein, refers to defined as a glycoprotein produced by B cells and plasma cells that binds with high specificity to an antigen (usually, but not always, a peptide) or a structurally similar antigen, that generated its production. Antibodies may be produced by any of the known methodologies and may be either polyclonal or monoclonal.

"Staining," as used herein, refers to any number of processes known to those in the field (typically utilizing dyes) that are used to visualize a specific component(s) and/or feature(s) of a cell or cells.

09709020.110300

"Alcohol," as used herein, refers to compounds that have hydroxyl functional groups bonded to saturated,  $sp^3$ -hybridized carbon atoms. The term "short-chain alcohol," as used herein, refers to alcohols that contain less than 10 carbon atoms. Examples of such short-chain alcohols comprise methanol, ethanol, propanol, butanol, pentanol, hexanol, heptanol, octanol, nonanol, including isomers thereof. The term "long-chain alcohol," as used herein, refers to fatty alcohols, in particular, the higher aliphatic, primary alcohols containing from 10 to 18 carbon atoms, preferably saturated and preferably straight-chain alcohol of the type obtainable by the industrial hydrogenation of native fatty acids. Typical representatives of the higher aliphatic alcohols for example the compounds n-dodecyl alcohol, n-tetradecyl alcohol, n-hexadecyl alcohol, n-octadecyl alcohol, n-octyl alcohol, n-decyl alcohol, undecyl alcohol, tridecyl alcohol.

"Sulfur donor," as used herein, refers to any sulfur-based compound that is capable of providing a sulfonic acid group in the formation of uridine-5-diphospho sulfoquinovose (UDP-SQ). Examples of such sulfur donors comprise sulfate, sulfite, sulfide, thiosulfate, sulfoglutathione, adenosine 5'-phosphosulfate (APS), and 3'-phosphoadenosine-5'-phosphosulfate (PAPS).

"Acid catalyst," as used herein, refers to any acidic compounds including the so-called Lewis acids, which catalyze the acetalization reaction between fatty alcohol and a sugar molecule. Examples of acids used for this purpose in industrial processes comprise mineral acids such as  $H_2SO_4$ ,  $HCl$ ,  $H_3PO_4$  or  $BF_3$ , or sulfonic acids or their salts. Examples of sulfonic acids comprise ortho-, meta- and para-toluenesulfonic acids, alkylbenzenesulfonic acids, secondary alkyl-sulfonic acids, sulfonic resins, alkylsulfates, alkylbenzenesulfonates, alkyl-sulfonates and sulfosuccinic acid.

"Alkyl sulfoquinovoside," as used herein, refers to a group of substances consisting of a glycosidic unit sulfonated at the C-6 position and acetalized at the C-1 position with an alcohol. In the context of the invention, alkyl sulfoquinovosides are understood to be the reaction products of UDP-sulfoquinovose and fatty alcohols. In its broadest sense, the term "alkyl" in alkyl sulfoquinovosides is intended to encompass the residue of an aliphatic C8-C18 alcohol, obtainable from natural fats, *i.e.* saturated

and unsaturated residues and also mixtures thereof, including those having different chain lengths. The terms alkyl oligosulfoquinovosides, alkyl polysulfoquinovosides apply to alkylated sulfoquinovosides of the type in which one alkyl residue in the form of the acetal is attached to more than one sulfoquinovoside residue, i.e. to a polysulfoquinovoside or oligosulfoquinovoside residue; these terms are regarded as synonymous with one another. Accordingly, alkyl monosulfoquinovoside is the acetal of a monosulfoquinovoside. Since the reaction products of the sugars and the fatty alcohols are generally mixtures, the term alkyl sulfoquinovoside is intended to encompass both alkyl monosulfoquinovosides and also alkyl poly(oligo) sulfoquinovosides.

"Nucleic acid sequence," "nucleotide sequence," and "polynucleotide sequence" as used herein refer to an oligonucleotide or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single-, or double-stranded, and represent the sense or antisense strand.

As used herein, the terms "oligonucleotides" and "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 100 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides, which can be used as a probe or amplifier.

The term "nucleotide sequence of interest" refers to any nucleotide sequence, the manipulation of which may be deemed desirable for any reason, by one of ordinary skill in the art. Such nucleotide sequences include, but are not limited to, coding sequences of structural genes (*e.g.*, enzyme-encoding genes, reporter genes, selection marker genes, oncogenes, drug resistance genes, growth factors, *etc.*), and of non-coding regulatory sequences that do not encode an mRNA or protein product (*e.g.*, promoter sequence, enhancer sequence, polyadenylation sequence, termination sequence, *etc.*).

"Amino acid sequence," "polypeptide sequence," "peptide sequence," and "peptide" are used interchangeably herein to refer to a sequence of amino acids.

The term "portion" when used in reference to a nucleotide sequence refers to fragments of that nucleotide sequence. The fragments may range in size from 5

nucleotide residues to the entire nucleotide sequence minus one nucleic acid residue. The term "portion" when used in reference to an amino acid sequence refers to fragments of the amino acid sequence. The fragments may range in size from 3 amino acids to the entire amino acid sequence minus one amino acid residue.

5           An oligonucleotide sequence which is a "homolog" of a first nucleotide sequence is defined herein as an oligonucleotide sequence which exhibits greater than or equal to 50% identity, and more preferably greater than or equal to 70% identity, to the first nucleotide sequence when sequences having a length of 10 bp or larger are compared.

10           DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotide is referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring. An end of an oligonucleotide is referred to as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of another mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends. In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements. This terminology reflects that transcription proceeds in a 5' to 3' direction along the DNA strand. The promoter and enhancer elements which direct transcription of a linked gene are generally located 5' or upstream of the coding region. However, enhancer elements can exert their effect even when located 3' of the promoter element and the coding region. Transcription termination and polyadenylation signals are located 3' or downstream of the coding region.

25           The term "cloning" as used herein, refers to the process of isolating a nucleotide sequence from a nucleotide library, cell or organism for replication by recombinant techniques.

The term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

The term "recombinant protein" or "recombinant polypeptide" as used herein refers to a protein molecule which is expressed using a recombinant DNA molecule.

As used herein, the terms "vector" and "vehicle" are used interchangeably in reference to nucleic acid molecules that transfer DNA segment(s) from one cell to another.

As used herein, the terms "complementary" or "complementarity" are used in reference to "polynucleotides" and "oligonucleotides" (which are interchangeable terms that refer to a sequence of nucleotides) related by the base-pairing rules. For example, the sequence "5'-CAGT-3'," is complementary to the sequence "5'-ACTG-3'." Complementarity can be "partial" or "total." "Partial" complementarity is where one or more nucleic acid bases is not matched according to the base pairing rules. "Total" or "complete" complementarity between nucleic acids is where each and every nucleic acid base is matched with another base under the base pairing rules. The degree of complementarity between nucleic acid strands may have significant effects on the efficiency and strength of hybridization between nucleic acid strands. This may be of particular importance in amplification reactions, as well as detection methods which depend upon binding between nucleic acids.

The terms "homology" and "homologous" as used herein in reference to nucleotide sequences refer to a degree of complementarity with other nucleotide sequences. There may be partial homology or complete homology (*i.e.*, identity). A nucleotide sequence which is partially complementary (*i.e.*, "substantially homologous") to a nucleic acid sequence is one that at least partially inhibits a completely complementary sequence from hybridizing to a target nucleic acid sequence. The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding

(i.e., the hybridization) of a completely homologous sequence to a target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding the probe will not hybridize to the second non-complementary target.

As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. "Stringency" typically occurs in a range from about  $T_m$ °C to about 20°C to 25°C below  $T_m$ . As will be understood by those of skill in the art, a stringent hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences. Under "stringent conditions" the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO: 5, and SEQ ID NO:6, or portions thereof, will hybridize to its exact complement and closely related sequences.

Low stringency conditions comprise conditions equivalent to binding or hybridization at 68°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$  and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1 % SDS, 5X Denhardt's reagent (50X Denhardt's contains per 500 ml: 5 g Ficoll (Type 400, Pharmacia), 5 g BSA (Fraction V; Sigma)) and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 2.0X SSPE, 0.1% SDS at room temperature when a probe of about 100 to about 1000 nucleotides in length is employed.

It is well known in the art that numerous equivalent conditions may be employed to comprise low stringency conditions; factors such as the length and nature (DNA, RNA, base composition) of the probe and nature of the target (DNA, RNA, base composition, present in solution or immobilized, etc.) and the concentration of the



09706000 "1108000  
salts and other components (e.g., the presence or absence of formamide, dextran sulfate, polyethylene glycol), as well as components of the hybridization solution may be varied to generate conditions of low stringency hybridization different from, but equivalent to, the above listed conditions. In addition, conditions which promote hybridization under conditions of high stringency (e.g., increasing the temperature of the hybridization and/or wash steps, the use of formamide in the hybridization solution, etc.) are well known in the art. High stringency conditions, when used in reference to nucleic acid hybridization, comprise conditions equivalent to binding or hybridization at 68°C in a solution consisting of 5X SSPE, 1 % SDS, 5X Denhardt's reagent and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 0.1X SSPE and 0.1 % SDS at 68°C when a probe of about 100 to about 1000 nucleotides in length is employed.

When used in reference to a double-stranded nucleic acid sequence such as a cDNA or genomic clone, the term "substantially homologous" refers to any probe which can hybridize either partially or completely to either or both strands of the double-stranded nucleic acid sequence under conditions of low stringency as described above.

When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous" refers to any probe which can hybridize to the single-stranded nucleic acid sequence under conditions of low stringency as described above.

As used herein, the term "hybridization" is used in reference to the pairing of complementary nucleic acids using any process by which a strand of nucleic acid joins with a complementary strand through base pairing to form a hybridization complex. Hybridization and the strength of hybridization (i.e., the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementarity between the nucleic acids, stringency of the conditions involved, the  $T_m$  of the formed hybrid, and the G:C ratio within the nucleic acids.

As used herein the term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bounds between complementary G and C bases and between complementary A and T bases;

these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (*e.g.*,  $C_0t$  or  $R_0t$  analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized to a solid support (*e.g.*, a nylon membrane or a nitrocellulose filter as employed in Southern and Northern blotting, dot blotting or a glass slide as employed in *in situ* hybridization, including FISH (fluorescent *in situ* hybridization)).

As used herein, the term " $T_m$ " is used in reference to the "melting temperature." The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. The equation for calculating the  $T_m$  of nucleic acids is well known in the art. As indicated by standard references, a simple estimate of the  $T_m$  value may be calculated by the equation:  $T_m = 81.5 + 0.41(\% G + C)$ , when a nucleic acid is in aqueous solution at 1 M NaCl (*see e.g.*, Anderson and Young, Quantitative Filter Hybridization, in *Nucleic Acid Hybridization* [1985]). Other references include more sophisticated computations which take structural as well as sequence characteristics into account for the calculation of  $T_m$ .

"Amplification" is defined herein as the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction technologies well known in the art (*see, e.g.*, Dieffenbach and Dveksler, *PCR Primer, a Laboratory Manual*, Cold Spring Harbor Press, Plainview NY [1995]). As used herein, the term "polymerase chain reaction" ("PCR") refers to the methods of U.S. Patent Nos. 4,683,195, 4,683,202, and 4,965,188, all of which are hereby incorporated by reference, which describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. The length of the amplified segment of the desired target sequence is determined by the relative positions of two oligonucleotide primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" (hereinafter

"PCR"). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified."

With PCR, it is possible to amplify a single copy of a specific target sequence in genomic DNA to a level detectable by several different methodologies (*e.g.*, hybridization with a labeled probe; incorporation of biotinylated primers followed by avidin-enzyme conjugate detection; incorporation of <sup>32</sup>P-labeled deoxynucleotide triphosphates, such as dCTP or dATP, into the amplified segment). In addition to genomic DNA, any oligonucleotide sequence can be amplified with the appropriate set of primer molecules. In particular, the amplified segments created by the PCR process itself are, themselves, efficient templates for subsequent PCR amplifications.

The terms "reverse transcription polymerase chain reaction" and "RT-PCR" refer to a method for reverse transcription of an RNA sequence to generate a mixture of cDNA sequences, followed by increasing the concentration of a desired segment of the transcribed cDNA sequences in the mixture without cloning or purification. Typically, RNA is reverse transcribed using a single primer (*e.g.*, an oligo-dT primer) prior to PCR amplification of the desired segment of the transcribed DNA using two primers.

As used herein, the term "primer" refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, (*i.e.*, in the presence of nucleotides and of an inducing agent such as DNA polymerase and at a suitable temperature and pH). The primer is preferably single stranded for maximum efficiency in amplification, but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent. The exact

lengths of the primers will depend on many factors, including temperature, source of primer and the use of the method.

As used herein, the term "probe" refers to an oligonucleotide (*i.e.*, a sequence of nucleotides), whether occurring naturally as in a purified restriction digest or produced synthetically, recombinantly or by PCR amplification, which is capable of hybridizing to another oligonucleotide of interest. A probe may be single-stranded or double-stranded. Probes are useful in the detection, identification and isolation of particular gene sequences. It is contemplated that any probe used in the present invention will be labeled with any "reporter molecule," so that it is detectable in any detection system, including, but not limited to enzyme (*e.g.*, ELISA, as well as enzyme-based histochemical assays), fluorescent, radioactive, and luminescent systems. It is not intended that the present invention be limited to any particular detection system or label.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double- or single-stranded DNA at or near a specific nucleotide sequence.

As used herein, the term "an oligonucleotide having a nucleotide sequence encoding a gene" means a nucleic acid sequence comprising the coding region of a gene, *i.e.* the nucleic acid sequence which encodes a gene product. The coding region may be present in either a cDNA, genomic DNA or RNA form. When present in a DNA form, the oligonucleotide may be single-stranded (*i.e.*, the sense strand) or double-stranded. Suitable control elements such as enhancers, promoters, splice junctions, polyadenylation signals, *etc.* may be placed in close proximity to the coding region of the gene if needed to permit proper initiation of transcription and/or correct processing of the primary RNA transcript. Alternatively, the coding region utilized in the expression vectors of the present invention may contain endogenous enhancers, splice junctions, intervening sequences, polyadenylation signals, *etc.* or a combination of both endogenous and exogenous control elements.

The term "promoter," "promoter element," or "promoter sequence" as used herein, refers to a DNA sequence which when placed at the 5' end of (*i.e.*, precedes) an oligonucleotide sequence is capable of controlling the transcription of the oligonucleotide sequence into mRNA. A promoter is typically located 5' (*i.e.*, upstream) of an oligonucleotide sequence whose transcription into mRNA it controls, and provides a site for specific binding by RNA polymerase and for initiation of transcription.

As used herein, the terms "nucleic acid molecule encoding," "nucleotide encoding," "DNA sequence encoding," and "DNA encoding" refer to the order or sequence of deoxyribonucleotides along a strand of deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of amino acids along the polypeptide (protein) chain. The DNA sequence thus codes for the amino acid sequence.

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" refers to a nucleic acid sequence that is separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is nucleic acid present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids are nucleic acids such as DNA and RNA which are found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. However, isolated nucleic acid encoding a polypeptide of interest includes, by way of example, such nucleic acid in cells ordinarily expressing the polypeptide of interest where the nucleic acid is in a chromosomal or extrachromosomal location different from that of natural cells, or is otherwise flanked by a different nucleic acid sequence than that found in nature. The isolated nucleic acid or oligonucleotide may be present in single-stranded or double-stranded form. Isolated nucleic acid can be readily identified (if desired) by a variety of techniques (*e.g.*, hybridization, dot blotting, etc.). When an isolated nucleic acid or

20

25

30

oligonucleotide is to be utilized to express a protein, the oligonucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide may be single-stranded). Alternatively, it may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide may be double-stranded).

5 As used herein the term "coding region" when used in reference to a structural gene refers to the nucleotide sequences which encode the amino acids found in the nascent polypeptide as a result of translation of a mRNA molecule. The coding region is bounded, in eukaryotes, on the 5' side by the nucleotide triplet "ATG" which encodes the initiator methionine and on the 3' side by one of the three triplets which  
10 specify stop codons (*i.e.*, TAA, TAG, TGA).

As used herein, the term "gene" means the deoxyribonucleotide sequences comprising the coding region of a structural gene. A "gene" may also include non-translated sequences located adjacent to the coding region on both the 5' and 3' ends such that the gene corresponds to the length of the full-length mRNA. The sequences  
15 which are located 5' of the coding region and which are present on the mRNA are referred to as 5' non-translated sequences. The sequences which are located 3' or downstream of the coding region and which are present on the mRNA are referred to as 3' non-translated sequences. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding  
20 region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are segments of a gene which are transcribed into heterogenous nuclear RNA (hnRNA); introns may contain regulatory elements such as enhancers. Introns are removed or "spliced out" from the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The  
25 mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

In addition to containing introns, genomic forms of a gene may also include sequences located on both the 5' and 3' end of the sequences which are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions  
30 (these flanking sequences are located 5' or 3' to the non-translated sequences present

on the mRNA transcript). The 5' flanking region may contain regulatory sequences such as promoters and enhancers which control or influence the transcription of the gene. The 3' flanking region may contain sequences which direct the termination of transcription, post-transcriptional cleavage and polyadenylation.

5           The term "transgenic" when used in reference to a cell refers to a cell which contains a transgene, or whose genome has been altered by the introduction of a transgene. The term "transgenic" when used in reference to a tissue or to a plant refers to a tissue or plant, respectively, which comprises one or more cells that contain a transgene, or whose genome has been altered by the introduction of a transgene.

10          Transgenic cells, tissues and plants may be produced by several methods including the introduction of a "transgene" comprising nucleic acid (usually DNA) into a target cell or integration of the transgene into a chromosome of a target cell by way of human intervention, such as by the methods described herein.

15           The term "transgene" as used herein refers to any nucleic acid sequence which is introduced into the genome of a cell by experimental manipulations. A transgene may be an "endogenous DNA sequence," or a "heterologous DNA sequence" (*i.e.*, "foreign DNA"). The term "endogenous DNA sequence" refers to a nucleotide sequence which is naturally found in the cell into which it is introduced so long as it does not contain some modification (*e.g.*, a point mutation, the presence of a selectable marker gene, *etc.*) relative to the naturally-occurring sequence. The term  
20          "heterologous DNA sequence" refers to a nucleotide sequence which is ligated to, or is manipulated to become ligated to, a nucleic acid sequence to which it is not ligated in nature, or to which it is ligated at a different location in nature. Heterologous DNA is not endogenous to the cell into which it is introduced, but has been obtained from  
25          another cell. Heterologous DNA also includes an endogenous DNA sequence which contains some modification. Generally, although not necessarily, heterologous DNA encodes RNA and proteins that are not normally produced by the cell into which it is expressed. Examples of heterologous DNA include reporter genes, transcriptional and

translational regulatory sequences, selectable marker proteins (*e.g.*, proteins which confer drug resistance), *etc.*

The term "foreign gene" refers to any nucleic acid (*e.g.*, gene sequence) which is introduced into the genome of a cell by experimental manipulations and may include gene sequences found in that cell so long as the introduced gene contains some modification (*e.g.*, a point mutation, the presence of a selectable marker gene, *etc.*) relative to the naturally-occurring gene.

The term "transformation" as used herein refers to the introduction of a transgene into a cell. Transformation of a cell may be stable or transient. The term "transient transformation" or "transiently transformed" refers to the introduction of one or more transgenes into a cell in the absence of integration of the transgene into the host cell's genome. Transient transformation may be detected by, for example, enzyme-linked immunosorbent assay (ELISA) which detects the presence of a polypeptide encoded by one or more of the transgenes. Alternatively, transient transformation may be detected by detecting the activity of the protein (*e.g.*,  $\beta$ -glucuronidase) encoded by the transgene (*e.g.*, the *uid A* gene) as demonstrated herein [*e.g.*, histochemical assay of GUS enzyme activity by staining with X-gluc which gives a blue precipitate in the presence of the GUS enzyme; and a chemiluminescent assay of GUS enzyme activity using the GUS-Light kit (Tropix)]. The term "transient transformant" refers to a cell which has transiently incorporated one or more transgenes. In contrast, the term "stable transformation" or "stably transformed" refers to the introduction and integration of one or more transgenes into the genome of a cell. Stable transformation of a cell may be detected by Southern blot hybridization of genomic DNA of the cell with nucleic acid sequences which are capable of binding to one or more of the transgenes. Alternatively, stable transformation of a cell may also be detected by the polymerase chain reaction of genomic DNA of the cell to amplify transgene sequences. The term "stable transformant" refers to a cell which has stably integrated one or more transgenes into the genomic DNA. Thus, a stable transformant is distinguished from a transient transformant in that, whereas genomic DNA from the



stable transformant contains one or more transgenes, genomic DNA from the transient transformant does not contain a transgene.

A "transformed cell" is a cell or cell line that has acquired the ability to grow in cell culture for many multiple generations, the ability to grow in soft agar and the ability to not have cell growth inhibited by cell-to-cell contact. In this regard, transformation refers to the introduction of foreign genetic material into a cell or organism. Transformation may be accomplished by any method known which permits the successful introduction of nucleic acids into cells and which results in the expression of the introduced nucleic acid. "Transformation" methods include, but are not limited to, such methods as microinjection, electroporation, and DNA particle "bombardment." Transformation may be accomplished through use of any expression vector. For example, the use of *Agrobacterium tumefaciens* to introduce foreign nucleic acid into plant cells is contemplated. Additionally, transformation refers to cells that have been transformed naturally, usually through genetic mutation.

The term "*Agrobacterium*" refers to a soil-borne, Gram-negative, rod-shaped phytopathogenic bacterium which causes crown gall. The term "*Agrobacterium*" includes, but is not limited to, the strains *Agrobacterium tumefaciens*, (which typically causes crown gall in infected plants), and *Agrobacterium rhizogens* (which causes hairy root disease in infected host plants). Infection of a plant cell with *Agrobacterium* generally results in the production of opines (e.g., nopaline, agropine, octopine, etc.) by the infected cell. Thus, *Agrobacterium* strains which cause production of nopaline (e.g.' strain LBA4301, C58, A208) are referred to as "nopaline-type" *Agrobacteria*; *Agrobacterium* strains which cause production of octopine (e.g.' strain LBA4404, Ach5, B6) are referred to as "octopine-type" *Agrobacteria*; and *Agrobacterium* strains which cause production of agropine (e.g., strain EHA105, EHA101, A281) are referred to as "agropine-type" *Agrobacteria*.

The terms "bombarding," "bombardment," and "biolistic bombardment" refer to the process of accelerating particles towards a target biological sample (e.g., cell, tissue, etc.) to effect wounding of the cell membrane of a cell in the target biological

sample and/or entry of the particles into the target biological sample. Methods for biolistic bombardment are known in the art (e.g., U.S. Patent No. 5,584,807, the contents of which are herein incorporated by reference), and are commercially available (e.g., the helium gas-driven microprojectile accelerator (PDS-1000/He) (BioRad)).

The term "microwounding" when made in reference to plant tissue refers to the introduction of microscopic wounds in that tissue. Microwounding may be achieved by, for example, particle bombardment as described herein.

The term "plant" as used herein refers to a plurality of plant cells which are largely differentiated into a structure that is present at any stage of a plant's development. Such structures include, but are not limited to, a fruit, shoot, stem, leaf, flower petal, etc. The term "plant tissue" includes differentiated and undifferentiated tissues of plants including, but not limited to, roots, shoots, leaves, pollen, seeds, tumor tissue and various types of cells in culture (e.g., single cells, protoplasts, embryos, callus, protocorm-like bodies, etc.). Plant tissue may be *in planta*, in organ culture, tissue culture, or cell culture.

The term "embryonic cell" as used herein in reference to a plant cell refers to one or more plant cells (whether differentiated or un-differentiated) which are capable of differentiation into a plant tissue or plant. Embryonic cells include, without limitation, protoplasts such as those derived from the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Daucus*, *Arabidopsis*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Hyoscyamus*, *Lycopersicon*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Hererocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browaalia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, and *Datura*. Also included are embryos (such as those from sorghum, maize, banana), embryonic meristems (such as those from soybean), embryogenic callus (such as from sugarcane), protocorm-like bodies (such as from pineapple), and embryogenic cells as exemplified

by those from garlic. The ability of an embryonic cell to differentiate into a plant is determined using methods known in the art. For example, differentiation of pineapple protocorm-like bodies into shoots may be accomplished by culturing the protocorm-like body on agar-solidified hormone-free modified Murashige & Skoog (MS) medium or on agar-solidified PM2 medium (U.S. Patent No. 6,091,003 incorporated by reference). Differentiation into pineapple roots may be accomplished by culture of protocorm-like bodies in liquid modified MS medium containing 1 mg/L NAA.

The term "conjugation" as used herein refers to the process in which genetic material is transferred from one microorganism to another involving a physical connection or union between the two cells. This process is commonly known to occur in bacteria, protozoa, and certain algae and fungi.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to methods for the synthesis and subsequent modification of uridine-5'-diphospho sulfoquinovose (UDP-SQ).

### 1. Method for biosynthesis of uridine-5'-diphospho sulfoquinovose (UDP-SQ)

The methods of the present invention comprise the utilization of recombinant enzymes from *Arabidopsis thaliana*, UDP-glucose, and a sulfur donor to synthesize UDP-SQ. Although the present invention is not limited by any specific reaction mechanism, in one embodiment, the production of UDP-SQ from a reaction mixture comprising UDP-glucose, *Arabidopsis thaliana* recombinant SQD1 enzyme protein, and sulfite is contemplated (See Figure 1).

### Biosynthesis of uridine-5'-diphosphosulfoquinovose (UDP-SQ)

*Arabidopsis thaliana* recombinant SQD1 enzyme protein catalyzes the formation of the sulfonic acid precursor, UDP-SQ, from UDP-glucose and a sulfur donor. In one embodiment, the UDP-SQ production reaction is carried out in a buffer comprising purified SQD1 protein, Na<sub>2</sub>SO<sub>3</sub>, radiolabeled UDP-glucose, and Tris for 40 minutes at 37°C. The reaction mixture is then heat denatured to inactivate the recombinant enzyme, and centrifuged at 10,000 X g for 5 minutes. The production of UDP-SQ as a reflection of SQD1 activity is detected as described below.

The biosynthesis of UDP-SQ as contemplated by the present invention is not limited to any specific pH value. In one embodiment, the pH is between 7.0 and 9.5. In a preferred embodiment, the pH of the reaction is 7.5.

Although the present invention is not limited to employing any specific sulfur donor, in one embodiment, the sulfur donor is selected from the group comprising sulfate, sulfide, thiosulfate, sulfoglutathione, adenosine 5'-phosphosulfate (APS), and 3'-phosphoadenosine-5'-phosphosulfate (PAPS). In a preferred embodiment, the sulfur donor is sulfite. (See also, Example 1).

**Detection of uridine-5'-diphosphosulfoquinovose (UDP-SQ) produced by the method of the present invention**

The present invention is not limited by any specific means of detecting UDP-SQ as the end product of the method of biosynthesis described above. In one embodiment, the means for detecting the production of UDP-SQ comprises using high performance liquid chromatography (HPLC) is as follows. For example, the heat denatured reaction mixture is subjected to analysis by HPLC (Waters Corp., Milford, MA) employing a Beckman (Fullerton, CA) Ultrasphere ODS column (4.6 mm X 25 cm, particle size 5  $\mu$ M) kept at 42°C. Substrates and products are separated by applying a linear gradient of 30 mM  $\text{KH}_2\text{PO}_4$ , 2mM tetrabutylammonium hydroxide (Fisher Scientific, Fair Lawn, NJ), adjusted to pH 6.0 with KOH, to HPLC grade acetonitrile (EM Science, Gibbstown, NJ) with a flow rate of 1 ml per minute over 45 minutes. In the HPLC system described above, the major compound produced by the reaction co-chromatographed with authentic UDP-SQ, indicating that this compound was UDP-SQ, and that the purified SQD1 catalyzed the synthesis of the UDP-SQ produced in the assay

**Production of *Arabidopsis thaliana* recombinant SQD1 enzyme protein**

Essingman *et al.*, "Phosphate Availability Affects the Thylakoid Lipid Composition and Expression of SQD1, a Gene Required for Sulfolipid Biosynthesis in *Arabidopsis thaliana*," *Proc. Natl. Acad. Sci. USA*, 95: 1950-955 (1998) discloses the production of *A. thaliana* recombinant SQD1 protein in *Escherichia coli* using a PCR-based strategy, and speculates that SQD1 is involved in the biosynthesis of UDP-SQ from UDP-Glucose.

The present invention is not limited by any particular method for the production of the recombinant SQD1 enzyme used in the production of UDP-SQ. In one embodiment, a means for the production of *Arabidopsis thaliana* recombinant SQD1 enzyme protein, having the amino acid/nucleic acid sequence noted in SEQ ID NO:5, is as follows.

In order to isolate *A. thaliana* genes encoding enzymes involved in the head group biosynthesis of thylakoid membranes, the dbEST database of expressed sequence tags was searched with the predicted amino acid sequence of the bacterial *sqdB* genes using TBLASTN. Through said search, a partial rice cDNA (EST D46477) was found that encodes a putative protein with high sequence similarity to the bacterial *sqdB* gene product. (See Figure 18: SEQ ID NO: 9). A fragment of the partial rice cDNA was used as a probe to screen a *A. thaliana* PRL2 cDNA library by heterologous DNA hybridization. Although the present invention is not limited by any specific hybridization conditions or membranes, in one embodiment, Hybond N+ (Amersham) membranes were used, and hybridization was performed at 53°C in sodium phosphate buffer (pH 7.2) containing SDS, EDTA, and BSA. After hybridization, the membrane was washed twice for 20 minutes in a SSPE, SDS solution at 53°C. Several cDNA clones were isolated, including one with an insert of 1,799 base-pairs, which was sequenced (GenBank accession No. AF022082)(See Figure 14: SEQ ID NO: 6). The corresponding locus of *A. thaliana* was designated SQD1 and the plasmid containing the cDNA with the 1,799 bp insert was designated pSQD1. (See also, Example 2.a.).

The present invention is not limited to any specific means of expressing recombinant SQD1 protein. In one embodiment, in order to express recombinant SQD1 protein in *Escherichia coli*, a fragment of pSQD1 was cloned into the His-tag expression vector, pQE-30 (QIAGEN, Inc., Valencia, CA: Cat.# 32149)(See Figure 3) using a PCR-based strategy. The present invention is not limited to the use of any specific protein expression vector or system. In one embodiment, the protein expression vector is selected from the group comprising pQE-9, pQE-16, pQE-31, pQE-32, pQE-40, pQE-60, pQE-70, pQE-80, pQE-81, pQE-82, pQE-100 (all available from QIAGEN, Inc., Valencia, CA). In another embodiment, the protein expression vector is pACYC184 (New England Biolabs, Beverly, MA: Cat.# E4152S). (See Figure 6). In a preferred embodiment, the protein expression vector is pQE-30. (See Figure 3).

09709030.140300

The present invention is not limited to any specific means of purifying recombinant SQD1 protein. In one embodiment, the resulting plasmid construct, pSQD1-TP, allowed the expression of the recombinant SQD1 protein in *E. coli* and the purification of the protein due to the selective binding of the six N-terminal histidine residues of the plasmid construct to nickel nitriloacetic acid (Ni-NTA) agarose resin following the manufacturer's instructions. (QIAGEN, Inc., Valencia, CA: Cat.# 30210). The recombinant protein was eluted and stored in a buffer comprising glycerol, NaCl, and NaH<sub>2</sub>PO<sub>4</sub> (pH 7.5) at -20°C. The SQD1 protein was estimated to be approximately 95% pure by SDS-PAGE gel analysis. (See Figure 4).

#### 10      **Assay for Measuring SQD1 Activity**

The present invention is not limited to any specific means of measuring the activity of recombinant SQD1 protein produced by the invention. In one embodiment, an enzyme assay was developed to measure the conversion of UDP-glucose to UDP-SQ as a reflection of SQD1 activity. Basic activity assays were carried out at 37°C in a reaction mixture containing purified SQD1 protein, Na<sub>2</sub>SO<sub>3</sub>, radiolabeled UDP-glucose, and Tris (pH 7.5) in a total volume of 100 µl for 40 minutes. The reaction mixture was incubated for 10 minutes, heat denatured, centrifuged, and analyzed by HPLC. Substrates and products were separated by applying a linear gradient of KH<sub>2</sub>PO<sub>4</sub>, tetrabutylammonium hydroxide (Fisher Scientific, Fair Lawn, NJ), adjusted to pH 6.0 with KOH, to HPLC grade acetonitrile (EM Science, Gibbstown, NJ).

Incubation of the SQD1 protein with labeled UDP-glucose as described above resulted in the formation of two compounds with unique retention times as compared to UDP-glucose as analyzed by HPLC. In the HPLC system described above, one compound co-chromatographed with authentic UDP-SQ, indicating that this compound was UDP-SQ, and that the purified SQD1 catalyzed the synthesis of the UDP-SQ produced in the assay. (See also, Examples 1 & 2.b.).

## 2. The Biosynthesis of 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG)

The methods of the present invention further comprise the subsequent modification of UDP-SQ to form compounds including, but not limited to, 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG). Although the present invention is not limited by any specific reaction mechanism, in one embodiment, the production of SQDG from a reaction mixture comprising UDP-SQ, diacylglycerol, and a recombinant peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol, is contemplated as follows. (See Figure 5).

In one embodiment, SQDG is produced in a reaction containing means by reacting 100  $\mu$ M UDP-SQ, 100  $\mu$ M diacylglycerol, and 10  $\mu$ g of a substantially purified peptide that is a gene product encoded by the nucleic acid sequence set forth in SEQ ID NO: 1, in a 100  $\mu$ l reaction volume at 37°C for 40 minutes. In another embodiment, said peptide is a gene product encoded by the nucleic acid sequence set forth in SEQ ID NO: 3. In another embodiment, said peptide is a gene product encoded by the nucleic acid sequence set forth in SEQ ID NO: 4. In a further embodiment, said peptide is a gene product encoded by the nucleic acid sequence set forth in SEQ ID NO: 5.

The present invention is not limited by a specific means for verifying the production of SQDG by the method described above. In one embodiment, the production of SQDG is verified by the following means. Aliquots of the above reaction are analyzed by thin layer chromatography (TLC) on activated ammonium sulfate impregnated silica gel TLC plates with a solvent system containing acetone-toluene-water (91:30:8, vol/vol/vol). Products of the above reaction are then visualized with iodine vapor and identified by co-chromatography with an *Arabidopsis thaliana* leaf lipid extract known to contain SQDG. (See Figure 9). In another embodiment, production of SQDG is verified by quantitative analysis wherein reaction products are isolated from the TLC plates and used to prepare fatty acid methyl esters. The methyl esters are quantified by gas chromatography using myristic acid as the internal standard as described below.



## Detection of SQDG Production by Thin Layer Chromatography (TLC)

Randomly chosen colonies from a mutagenized population of *R. sphaeroides* cells known to produce the lipid, SQDG, are streaked as small patches (0.5 by 0.5 cm) on fresh Z-broth plates. Lipids are isolated from these patches by collecting cells onto the wide end of a flat toothpick and swirling the material in 75  $\mu$ l of chloroform-methanol (1:1, vol/vol) contained in polypropylene microcentrifuge tubes. Following the addition of 25  $\mu$ l of 1 N KCl-0.2 M  $H_3PO_4$ , the tubes are vortexed and centrifuged to separate the organic and aqueous phases. A 10  $\mu$ l aliquot is withdrawn from the lipid-containing lower phase and directly spotted onto an activated ammonium sulfate-impregnated silica gel thin layer chromatography (TLC) plate. For this purpose, Baker Si250 silica plates with a preadsorbent layer are prepared by soaking in 0.15 M ammonium sulfate for 30 seconds followed by air drying to complete dryness. Immediately prior to use, the plates are activated for 2.5 h at 120°C. Activation of ammonium sulfate-treated plates at 120°C produces sulfuric acid, which protonates phosphatidylglycerol, making it less polar. An acetone-benzene-water mixture (91:30:8, vol/vol/ vol) is employed as the solvent system. Lipids were visualized by spraying the plates with 50% sulfuric acid followed by heating at 160°C for 10 to 15 min to char the lipids.

## Quantitative Lipid Analysis to Verify the Production of SQDG

For each strain, three 50-ml cultures were grown in Sistrom's medium aerobically with shaking at 32°C in the dark. The cells are centrifuged, suspended in 0.5 ml of water, and extracted by vortexing with 4 ml of chloroform-methanol (1:1, vol/vol). Addition of 1.3 ml of 1 M KCl-0.2 M  $H_3PO_4$ , vortexing, and centrifugation results in phase partitioning of the lipids into the lower chloroform phase. The chloroform phase is removed and concentrated to 0.2 ml by evaporation under a stream of  $N_2$ . The sample is split, and the material is spotted onto activated (30 min at 110°C) silica TLC plates (Si250; Baker). The plates are developed in two dimensions, first with chloroform-methanol-water (65:25:4, vol/ vol/vol) and then with chloroform-acetone-methanol-acetic acid-water (50:20:10:10:5, by volume).

Lipids are visualized with iodine vapor, and after desorption of iodine, the spots were individually scraped into 8-ml screw-cap tubes. To the samples, 5 µg of myristic acid methyl ester in 0.1 ml of hexane was added as an internal standard, since only negligible amounts of endogenous myristic acid were found in the bacterial lipids. Fatty acid methyl esters are prepared by addition of 1 ml of anhydrous 1 N methanolic HCl (Supelco) followed by incubation at 80°C for 1 h. Following the addition of 1 ml of 0.95% (wt/vol) KCl, the fatty acid methyl esters were extracted into 1 ml of hexane and then dried to a volume of 0.1 ml.

Samples (2 µl each) are injected onto a gas chromatograph (Varian 2000) which was equipped with a 2.4-m column (2-mm inner diameter) packed with 3% SP-2310 and 2% SP-2300 on 100/120 Chromosorb WAW (Supelco). The carrier gas (N<sub>2</sub>) flow rate was adjusted to 20 ml/min, and the column temperature was set for 2 min at 180°C, increasing to 200°C over 10 min, and 4 min at 200°C. The fatty acid methyl esters were detected by a flame ionization detector, and the data were integrated by a Spectra Physics integrator. To calculate the relative amounts of the eight polar lipids included in the analysis, the amount of fatty acids contained in each lipid was calculated. The validity of calculation was based on the assumption that each of the lipids, including the unidentified lipids, contained two fatty acids per molecule and that the different lipids had a similar fatty acid composition.

## **Production and purification of a recombinant peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol**

### **a. Cyanobacterial Peptide**

The invention is not limited to a specific means for the expression of a recombinant peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol. In one embodiment, a means for the production of a substantially purified peptide encoded by the nucleic acid sequence as set forth in SEQ ID NO: 1, is as follows.

In one embodiment, in order to express recombinant SQDX protein in *Escherichia coli*, a 1,133 base-pair fragment of pSYB (See Example 4) containing

nucleotide numbers 1800-2933 of SEQ ID NO: 1 (GenBank Accession No. AF155063) was cloned into the His-tag expression vector, pQE-30 (QIAGEN, Inc., Valencia, CA: Cat.# 32149) using a PCR-based strategy. For this purpose, a forward primer having the nucleotide sequence 5'-TTT GGA TCC CGC ATC GCT CTC TTT-3' (SEQ ID NO: 12), and a reverse primer having the nucleotide sequence 5'-ATA AGC TTC GAG CTC AGG CCG CT-3' (SEQ ID NO: 13), were used such that *Bam*HI and *Hind*III sites were provided for cloning into pQE-30. The forward primer amplifies the beginning of the gene with the omission of the *Met* start site (ATG) and immediately starts at the second amino acid (See Figure 17: SEQ ID NO: 2). The reverse primer includes the stop codon of the *sqdX* gene in the resulting PCR product.

The present invention is not limited to the use of any specific protein expression vector or system. In one embodiment, the protein expression vector is selected from the group comprising pQE-9, pQE-16, pQE-31, pQE-32, pQE-40, pQE-60, pQE-70, pQE-80, pQE-81, pQE-82, pQE-100 (all available from QIAGEN, Inc., Valencia, CA). In another embodiment, the protein expression vector is pACYC184 (New England Biolabs, Beverly, MA: Cat.# E4152S). (See Figure 6). In a preferred embodiment, the protein expression vector is pQE-30. (See Figure 3).

The present invention is not limited to any specific means of purifying recombinant SQDX protein. In one embodiment, the resulting plasmid construct allowed the expression of the recombinant SQDX protein in *E. coli* and the purification of the protein due to the selective binding of the six N-terminal histidine residues of the plasmid construct to Ni-NTA agarose following the manufacturer's instructions. (QIAGEN, Inc., Valencia, CA: Cat.# 30210). The recombinant protein was eluted with 200 mM imidazole, which was subsequently removed by use of a Millipore Ultrafree 4 concentrator (Millipore, Inc., Bedford, MA). The protein was stored in 20% glycerol, 300 mM NaCl, and 25 mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.5) at -20°C.

**b. *Arabidopsis* Peptide - *Cyanobacteria sqdX* gene homologs**

In another embodiment, the production of a substantially purified, recombinant *Arabidopsis thaliana* peptide capable of transferring sulfoquinovose from UDP-SQ

onto diacylglycerol is contemplated. In one embodiment, a means for the production of *sqdX* gene homologs of *Arabidopsis thaliana* encoded by the nucleic acid sequences as set forth in SEQ ID NO: 3, SEQ ID NO: 4, and SEQ ID NO: 5 is described. A BLAST comparison of the Cyanobacterial *sqdX* gene to genomic sequence of *Arabidopsis thaliana* revealed several potential homologs. In one embodiment, *AtSQDX-1*, a homolog having 37% amino acid identity with the Cyanobacterial *sqdX* gene and a nucleic acid sequence as set forth in SEQ ID NO: 3 is contemplated. In another embodiment, *AtSQDX-2*, a homolog having 29% amino acid identity with the cyanobacterial *sqdX* gene is and a nucleic acid sequence as set forth in SEQ ID NO: 4 is contemplated. In a further embodiment, *AtSQDX-3*, a homolog having 32% amino acid identity with the Cyanobacterial *sqdX* gene and a nucleic acid sequence as set forth in SEQ ID NO: 5 is contemplated.

Although the present invention is not limited to the expression of any specific *Arabidopsis thaliana sqdX* homolog, in one embodiment, *AtSQDX-1* is cloned and expressed as follows.

Total RNA from leaves of two-week old *Arabidopsis* wild-type plants is isolated according to Logemann *et al.*, "Improved Method for the Isolation of RNA from Plant Tissues," *Anal. Biochem.*, 163: 16-20 (1987), as described below. In one embodiment, the *Arabidopsis* leaves are phosphate deprived to enrich for SQDX sequences. The isolated total RNA is then enriched for Poly A<sup>+</sup> mRNA using the Oligotex mRNA Mini Kit (QIAGEN Cat. No. 70022) following the manufacturer's instructions as described below. The mRNA is subjected to cDNA biosynthesis using the ProSTAR HF Single-Tube RT-PCR System (Stratagene, LaJolla, CA: Cat. No. 600164) following the manufacturer's instructions (as described below) in order to produce a cDNA containing the open reading frame of *AtSQDX-1*. Primers based on the available genomic sequence of *AtSQDX-1* (SEQ ID NO: 3)(GenBank Accession No. AL137189) are designed to allow in-frame cloning into the protein expression vector, pQE-30.

09709090 10900

In one embodiment, in order to express recombinant *AtSQDX-1* protein in *Escherichia coli*, a 1,410 base-pair fragment of pSYB comprising at least a portion of the nucleic acid sequence set forth in SEQ ID NO: 3 (GenBank Accession No. CAB69850) was cloned into the His-tag expression vector, pQE-30 (QIAGEN, Inc., Valencia, CA: Cat.# 32149) using a PCR-based strategy. For this purpose, a forward primer having the nucleotide sequence 5'-CGG GAT CCA TGA CGA CTC TTT CTT CTA TA-3' (SEQ ID NO: 14), and a reverse primer having the nucleotide sequence 5'-AAG GAT CCC TAC ACG TTA CCT TCC GGT A-3' (SEQ ID NO: 15), were used such that a *BamHI* site was provided for cloning into pQE-30.

The present invention is not limited with respect to any specific primers used to generate an *Arabidopsis thaliana sqdX* homolog. In another embodiment, the forward primer, 5'-AAG GAT CCA TGG CTT CAC AAA CCA AAC T-3' (SEQ ID NO: 16), and the reverse primer, 5'-GCG GAT CCT CAT ATT TTG AAA AAG CAC T-3' (SEQ ID NO: 17), produce the cDNA for the *AtSQDX-2* gene. In a further embodiment, the forward primer, 5'-AGG GTA CCA TGG AGG GAT TCG GTT ATC-3' (SEQ ID NO: 18), and the reverse primer, 5'-GCG GTA CCT TAA GGT CTA TGC ATT TGA C-3' (SEQ ID NO: 19), produce the cDNA for the *AtSQDX-3* gene.

The present invention is not limited to the cloning of any specific nucleotide sequence into a protein expression vector to produce a recombinant *A. thaliana* peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol. In one embodiment, a fragment of the *AtSQDX-2* gene comprising at least a portion of the nucleic acid sequence as set forth in SEQ ID NO: 4 is cloned into pQE-30. In another embodiment, a fragment of the *AtSQDX-3* gene comprising at least a portion of the nucleic acid sequence as set forth in SEQ ID NO: 5 is cloned into pQE-30.

The present invention is not limited to the use of any specific protein expression vector or system. In one embodiment, the protein expression vector is selected from the group pQE-9, pQE-16, pQE-31, pQE-32, pQE-40, pQE-60, pQE-70, pQE-80, pQE-81, pQE-82, pQE-100 (all available from QIAGEN, Inc., Valencia, CA).

09705030.10600

In another embodiment, the protein expression vector is pACYC184. (See Figure 6).  
In a preferred embodiment, the protein expression vector is pQE-30. (See Figure 3).

The present invention is not limited to any specific means of purifying a recombinant *Arabidopsis thaliana* peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol recombinant protein. In one embodiment, the resulting plasmid construct allowed the expression of the recombinant AtSQDX-1 protein in *E. coli* and the purification of the protein due to the selective binding of the six N-terminal histidine residues of the plasmid construct to Ni-NTA agarose resin following the manufacturer's instructions. (QIAGEN, Inc., Valencia, CA: Cat.# 30210). The recombinant protein was eluted with 200 mM imidazole, which was subsequently removed by use of a Millipore Ultrafree 4 concentrator (Millipore, Inc., Bedford, MA). The protein was stored in 20% glycerol, 300 mM NaCl, and 25 mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.5) at -20°C.

In another embodiment, purification of a AtSQDX-2 gene product capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol is contemplated. In a further embodiment, purification of a AtSQDX-3 gene product is contemplated.

#### **Isolation of Total RNA from *Arabidopsis thaliana* Tissues**

It is not intended that the invention be limited by any specific method to isolate total RNA from *A. thaliana* tissues. In one embodiment, total RNA is isolated from said tissues by guanidine hydrochloride extraction as follows. Said tissues are frozen in liquid nitrogen and homogenized to a fine powder using a Waring blender. For small amounts of tissue (less than 0.5 g), a rotating pin in a 1.5-ml Eppendorf tube is used to homogenize the tissue. The extract is homogenized further at room temperature by the addition of 2 volumes of a guanidine buffer comprising 8 M guanidine hydrochloride, 20 mM MES (4-morpholineethansulfonic acid), 20 mM EDTA, and 50 mM 2-mercaptoethanol at pH 7.0.

The guanidine hydrochloride extract is centrifuged in a precooled (4°C) centrifuge for 10 minutes at 10,000 rpm. Subsequently the RNA-containing

supernatant is filtered through one layer of cheesecloth to get rid of floating particles. At least 0.2-1.0 vol of phenol/chloroform/IAA is added to extract proteins. After extraction the mixture is centrifuged for 45 minutes at 10,000 rpm at room temperature to separate the phases. The RNA-containing aqueous phase is collected and mixed with precooled 0.7 volumes of ethanol and 0.2 volumes of 1 M acetic acid for precipitating the RNA and leaving DNA and residual proteins in the supernatant. An overnight incubation at -20°C, or a 1 hour incubation at -70°C, is recommended.

The precipitated RNA is pelleted at 10,000 rpm for 10 min and washed twice with sterile 3 M sodium acetate at pH 5.2 at room temperature. Low-molecular-weight RNAs and contaminating polysaccharides dissolve, whereas intact RNA stays as a pellet after centrifugation for 5 minutes at 10,000 rpm. The salt is removed by a final wash with 70% ethanol and the RNA pellet is subsequently dissolved in sterile water and stored at 20°C until needed.

#### **Poly A+ mRNA Isolation from *Arabidopsis thaliana* Total RNA**

The present invention is not limited to any specific means of isolating Poly A+ mRNA from the total RNA of *Arabidopsis thaliana* leaves. In one embodiment, Poly A+ mRNA was isolated from *A. thaliana* leaf total RNA with the Oligotex mRNA Mini Kit (QIAGEN Cat. No. 70022) following the manufacturer's instructions as follows.

The Oligotex Suspension is heated to 37°C in a heating block, mixed by vortexing, and placed at room temperature. A sample containing 0.25 mg of *A. thaliana* leaf total RNA is pipetted into an RNase-free 1.5-ml microcentrifuge tube, and the volume of the reaction is adjusted to 0.25 ml with RNase-free water. A volume of 0.25 ml of Buffer OBB and 0.015 ml of Oligotex Suspension are added to the reaction. The contents are mixed thoroughly by pipetting. The sample incubated for 3 minutes at 70°C in a water bath or heating block in order to disrupt secondary structure of the RNA. The sample is removed from the heating block, and placed at room temperature (20° to 30°C) for 10 minutes to allow hybridization between the

oligo dT30 of the Oligotex particle and the poly-A tail of the mRNA. The Oligotex:mRNA complex is pelleted by centrifugation for 2 minutes at maximum speed (14,000–18,000 x g), and the supernatant is removed by pipetting.

The Oligotex:mRNA pellet is resuspended in 400 µl Buffer OW2 by vortexing, and pipetted onto a small spin column supplied with the kit. The spin column is centrifuged for 1 minute at maximum speed (14,000–18,000 x g). The spin column is transferred to a new RNase-free 1.5-ml microcentrifuge tube, and 400 µl of Buffer OW2 is applied to the column. The spin column is centrifuged for 1 minute at maximum speed and the flow-through fraction is discarded.

The spin column is transferred to another 1.5-ml microcentrifuge tube. A volume of 20–100 µl hot (70°C) Buffer OEB is pipetted onto the column. The resin is resuspended by pipetting up and down three or four times to allow elution of the mRNA, and centrifuged for 1 minute at maximum speed to pellet the suspension. The flow-through fraction, which contains the Poly A+ mRNA isolated, is stored at -20°C until used.

#### **Biosynthesis of *Arabidopsis thaliana* cDNA**

Although the present invention is not limited to any specific method for the biosynthesis of *Arabidopsis thaliana* cDNA, in one embodiment, said cDNA was biosynthesized using the ProSTAR HF Single-Tube RT-PCR System (Stratagene, LaJolla, CA: Cat. No. 600164) as follows.

Control and experimental reactions are prepared by adding the following components to separate sterile 0.5-ml microcentrifuge tubes in order:

##### Control Reaction

40.5 µl of RNase-free water (not DEPC-treated water)

5.0 µl of 10× HF RT-PCR buffer

1.0 µl of control primer set (200 ng/µl)

1.0 µl of dNTP mix (40 mM)

1.0 µl of control mRNA



### Experimental Reaction

39.5 µl of RNase-free water (not DEPC-treated water)

5.0 µl of 10× HF RT-PCR buffer

1.0 µl of forward primer (100 ng)

5 1.0 µl of reverse primer (100 ng)

1.0 µl of dNTP mix (40 mM)

1.0 µl (0.1–10 ng) of isolated Poly A<sup>+</sup> mRNA.

Just before use, 0.5 µl of StrataScript RT (20 U/µl) is diluted to a 8.0 µl final volume with 6.7 µl of RNase-free water and 0.8 µl of 10× HF RT-PCR buffer. A volume of 1.0 µl of the diluted StrataScript RT is added to each reaction. A volume of 0.5 µl of TaqPlus Precision DNA polymerase mixture is then added to each reaction. The reaction is vortexed gently without creating bubbles. Both the control and experimental reactions are placed in a GeneAmp PCR System 9600 thermal cycler (Applied Biosystems, Foster City, CA: Cat.# N801-0001). The reaction is then subjected to the following thermal-cycling program to both synthesize first-strand cDNA from the mRNA template and to amplify the cDNA via PCR: 1 cycle at 42°C for 30 minutes; 1 cycle at 95°C for 1 minute; 40 cycles comprised of 95°C for 30 seconds, 60°C for 30 seconds, and 68°C for 2 minutes; and 1 cycle at 68°C for 10 minutes.

Upon completion of the thermal-cycling program, the RT-PCR products are analyzed by 1.0% (w/v) agarose gel electrophoresis. RT-PCR amplification of the control reaction, which contains the control mRNA and the control primer set, yields a 500 base pair product. The reaction products will be readily visible by UV transillumination of the ethidium bromide-stained agarose gel. The products containing the cDNA produced by the above reaction are stored at –20°C until needed.

### **Co-Expression of *Arabidopsis thaliana* Recombinant Peptides**

It is not intended that the invention be limited to the independent expression of a peptide capable of catalyzing the conversion of UDP-Glc and a sulfur donor to UDP-

SQ in a single host organism or plant. Moreover, it is also not intended that the invention be limited to the independent expression of a second peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol in a single host organism or plant. In one embodiment, the invention contemplates the co-expression of both of the peptides described above in a single host organism or plant. In one embodiment, co-expression of the peptides SQD1 and SQDX (for example, in separate protein expression vectors) in *E. coli*, such that the sulfolipid biosynthetic pathway is reconstituted, is contemplated as follows.

In order to express two proteins in *E. coli*, two compatible plasmids with the ability to express proteins, one for SQD1 and one for SQDX, are utilized. Each plasmid must have a different antibiotic resistance in order to select for transformants with the correct combination of plasmids. The plasmid pQE-30 provides ampicillin resistance, whereas the plasmid, pACYC184, provides chloramphenicol resistance. The SQD1 coding region, along with the protein expression cassette of pQE-30, is removed from this plasmid using the restriction enzymes *Xho* I and *Pvu* II, and ligated into the pACYC184 plasmid (New England Biolabs, Beverly, MA: Cat.# E4152S) (See Figure 6) cut with *Sal* I and *EcoR* V. The M15 cell line (QIAGEN, Inc., Valencia, CA) is transformed with a pQE-30/SQDX protein expression construct (as described above). The SQD1/pACYC184 expression construct is transformed into the M15 cell line containing the pQE-30/SQDX expression vector. Upon induction of expression with 1-5 mM isopropyl- $\beta$ -D-thiogalactoside (IPTG)(Amersham Pharmacia Biotech, Piscataway, NJ: Cat.# 27-3054-03), both proteins are expressed.

The present invention is not limited to the use of any specific protein expression vector to produce co-expression of the two recombinant peptides. In one embodiment, the protein expression vector is selected from the group comprising pBK-CMV (Stratagene, LaJolla, CA: Cat.# 212209), pGEX-6P-1 (Amersham Pharmacia Biotech, Piscataway, NJ: Cat.# 27-4597-01), or pUC19 (New England Biolabs, Beverly, MA: Cat.# N3041S).

## Expression of of *Arabidopsis thaliana* Recombinant Peptides in Transgenic Plants

Transfer and expression of transgenes in plant cells is now routine practice to those skilled in the art. It has become a major tool to carry out gene expression studies and to attempt to obtain improved plant varieties of agricultural or commercial interest. The present invention is not limited to the expression of a first peptide capable of catalyzing the conversion of UDP-Glc and a sulfur donor to UDP-SQ in a single host organism, or a second peptide capable of transferring sulfoquinovose from UDP-SQ onto diacylglycerol, in bacterial cells. The invention contemplates the expression of *Arabidopsis thaliana* recombinant peptides in transgenic plants as described by S. Clough and A. Bent, "Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*," *Plant J.*, 16: 735-43 (1998). (See Example 3).

In one embodiment, the general process for manipulating genes to be transferred into the genome of plant cells to result in the expression of a recombinant peptide is carried out in two phases. First, all the cloning and DNA modification steps are done in *E. coli*, and the plasmid containing the gene construct of interest is transferred by conjugation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria (*e.g.*, streptomycin), and the other that will express in plants (*e.g.*, a gene encoding for kanamycin resistance or a gene encoding for resistance to an herbicide such as hygromycin). Also present are restriction endonuclease sites for the addition of one or more transgenes operably linked to appropriate regulatory sequences and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the region that will be transferred to the plant.

In another embodiment, plant cells may be transformed by shooting into the cell, tungsten microprojectiles on which cloned DNA is precipitated. (See, e.g., Gordon-Kamm *et al.*, *Plant Cell*, 2: 603 (1990)). In one embodiment, the Biolistic Apparatus (Bio-Rad, Hercules, Calif.) is used for the shooting with a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast driving a plastic macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to go through. As a result, the plastic macroprojectile smashes against the stopping plate and the tungsten microprojectiles continue toward their target through the hole in the plate. For the present invention the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

It is not intended that the present invention be limited to the particular manner by which the expression of any specific recombinant *A. thaliana* peptide in plants is achieved. In one embodiment, a peptide encoded by the nucleic acid sequences as set forth in SEQ ID NO: 6 is expressed in plants. In another embodiment, a peptide encoded by the nucleic acid sequence as set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5 is expressed in plants. In a further embodiment, two recombinant *A. thaliana* peptides encoded by the group of nucleic acid sequences comprising SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, and SEQ ID NO: 6 are co-expressed in plants.

It is not intended that the present invention be limited by any particular plant cell type in which to generate the expression of *Arabidopsis thaliana* recombinant peptides. In one embodiment, the plant cell is derived from a monocotyledonous plant. In an alternative embodiment, the plant cell is derived from a dicotyledonous plant. In another embodiment, the plant cell is derived from a group comprising the genera Anacardium, Arachis, Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus,

Capsicum, Carthamus, Cocos, Coffea, Cucumis, Cucurbita, Daucus, Elaeis, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Lactuca, Linum, Lolium, Lupinus, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Olea, Oryza, Panieum, Pannesetum, Persea, Phaseolus, Pistachia, Pisum, Pyrus, Prunus, Raphanus, Ricinus, Secale, Senecio, Sinapis, Solanum, Sorghum, Theobromus, Trigonella, Triticum, Vicia, Vitis, Vigna, and Zea. In a preferred embodiment, the plant cell is derived from *Arabidopsis thaliana*.

### 3. Subsequent Modification of UDP-SQ to Produce Alkyl Sulfoquinovosides

The methods of the present invention further comprise the subsequent modification of UDP-SQ to form compounds including, but not limited to, alkyl sulfoquinovoside. (See Figure 7). The method of the invention is not limited to the production of alkyl sulfoquinovosides. Furthermore, the present invention is not limited by any specific reaction mechanism. In one embodiment, the present invention relates to a process for the production of alkyl sulfoquinovosides by reacting of a short-chain alcohol with sulfoquinovose resulting from the hydrolytic cleavage of UDP-sulfoquinovose in the presence of a suitable acid catalyst with elimination of water. The short-chain alkyl sulfoquinovoside is then transacetalized with a long-chain alcohol to form long-chain sulfoquinovosides.

Although the method of the invention is not limited with respect to the structure of the alkyl sulfoquinovoside produced, in one embodiment, alkyl sulfoquinovosides are a group of substances consisting of a glycosidic unit sulfonated at the C-6 position and acetalized at the C-1 position with an alcohol. In another embodiment, alkyl sulfoquinovosides are understood to be the reaction products of UDP-sulfoquinovose and fatty alcohols. In a preferred embodiment, the term "alkyl" in alkyl sulfoquinovosides is intended to encompass the residue of an aliphatic C8-C18 alcohol, preferably a fatty alcohol, obtainable from natural fats (*i.e.* saturated and unsaturated residues and also mixtures thereof, including those having different chain lengths).

09709090 110900

The terms alkyl oligosulfoquinovosides and alkyl polysulfoquinovosides apply to alkylated sulfoquinovosides in which one alkyl residue in the form of the acetal is attached to more than one sulfoquinovoside residue (*i.e.* to a polysulfoquinovoside or oligosulfoquinovoside residue). Accordingly, alkyl monosulfoquinovoside is the acetal of a monosulfoquinovoside. Since the reaction products of the sugars and the fatty alcohols are generally mixtures, the term alkyl sulfoquinovoside is intended to encompass both alkyl monosulfoquinovosides and also alkyl poly(oligo) sulfoquinovosides.

In one embodiment, the synthesis of alkyl sulfoquinovose is carried out by the transacetalization method with a short-chain alcohol. Although it is not intended that the methods recited by the present invention be limited to any particular short-chain alcohol, in one embodiment, the short-chain alcohol is selected from a group comprising methanol, ethanol, propanol, pentanol, hexanol, heptanol, octanol, nonanol, including isomers thereof. In a preferred embodiment, the short-chain alcohol is butanol.

In one embodiment, the synthesis of alkyl sulfoquinovosides starts from the hydrolytic cleavage of the UDP-Sulfoquinovose. Sulfoquinovose is then refluxed with an acidic catalyst in butanol, the water of the reaction is removed by distillation under vacuum. The purpose of the acid catalyst is to favor the reactions which involve the glucosidic bond.

Although it is not intended that the method of the invention be limited to any particular acid catalyst, in one embodiment, any acidic compound (including the so-called Lewis acids which catalyze the acetalization reaction between fatty alcohol and the sugar molecule) may be used as catalysts. In one embodiment, the acid catalyst is a mineral acids comprising  $H_2SO_4$ ,  $HCl$ ,  $H_3PO_4$  or  $BF_3$ . In another embodiment, the acid catalyst is a sulfonic acid or its salt, comprising ortho-toluenesulfonic acid, meta-toluenesulfonic acid, alkylbenzenesulfonic acid, secondary alkyl-sulfonic acid, sulfonic resin, alkylsulfate, alkylbenzenesulfonate, alkyl-sulfonate and sulfosuccinic acid. In a more preferred embodiment, the acid catalyst is para-toluenesulfonic acid.

Although it is not intended that the method of the invention be limited to a particular set of reaction conditions, in one embodiment, the reflux temperature is 118 °C; a vapor temperature of 95 to 110°C. is established with the formation of the lower boiling butanol/water mixture; the acetalization with the butanol is carried out under light vacuum (*i.e.* under a pressure of 800 to 950 mbar); and an azeotropic amount of butanol is removed with the water.

In one embodiment, the butyl sulfoquinovoside is subsequently treated under vacuum with the long-chain alcohol, in the presence of the acidic catalyst. In one embodiment, it is preferable to reduce the content of butyl sulfoquinovoside by removing butanol by distillation under reduced pressure of 10 mbar. In one embodiment, neutralization of the catalyst following the removal of butanol is preferably separated by an interim period of up to about 1 hour, under which the reaction mixture is stirred under normal pressure at temperatures of from 100 to 115 °C. In this manner, the reaction of the butyl sulfoquinovoside with the fatty alcohol can be continued under control.

Although it is not intended that the method of the invention be limited to a particular long-chain alcohol, in one embodiment, the long-chain alcohol is a fatty alcohol; more preferably, a higher aliphatic, primary alcohol containing from 8 to 18 carbon atoms; and even more preferably, a saturated and preferably straight-chain alcohol of the type obtainable by the industrial hydrogenation of native fatty acids. In one embodiment, the higher aliphatic alcohol is selected from a group comprising n-dodecyl alcohol, n-tetradecyl alcohol, n-octadecyl alcohol, n-octyl alcohol, n-decyl alcohol, undecyl alcohol, tridecyl alcohol. In another embodiment, the long-chain alcohol is a technical mixture of about 3 parts by weight lauryl alcohol and 1 part by weight myristyl alcohol. In another embodiment, the long-chain alcohol is a branched-chain primary alcohol including, but not limited to, oxoalcohol. In a preferred embodiment, the long-chain alcohol is n-hexadecyl alcohol.

Although it is not intended that the method of the invention be limited to a particular set of reaction conditions, in one embodiment, the reaction mixture comprising short and long chain alkyl sulfoquinovosides, a long chain alcohol, and an

acid catalyst, is subsequently cooled to a temperature below 95°C. In one embodiment, the acidic catalyst is subsequently neutralized by the addition of a base and the adjustment of the pH of the neutralized reaction mixture to a pH of at least 8. In a preferred embodiment, the pH of the neutralized reaction mixture is 8.5.

5           Although it is not intended that the method of the invention be limited to a particular base, in one embodiment, the base is selected from a group of organic or inorganic basic materials comprising the alkali metal bases such as alkali metal hydroxide, carbonates, and bicarbonates. In another embodiment, the base is selected from a group comprising the alkaline earth bases such as calcium oxide and  
10           magnesium oxide. In another embodiment, aluminum bases such as aluminum hydroxide or its basic alkali aluminum components are contemplated. In a further embodiment, the base is selected from a group comprising ammonia-based compounds, such as ammonium hydroxide, and amines including, but not limited to, primary, secondary tertiary and heterocyclic amines.

15           Although it is not intended that the method of the invention be limited to a particular temperature range for filtering the reaction mixture, in one embodiment, the reaction mixture is filtered at a temperature of from 80 to 90°C, and the excess fatty alcohol is removed by distillation at to a level below 5% by weight. In one  
20           embodiment, the sump temperature must be kept at levels at which the alkyl sulfoquinovoside is thermally stable. In a preferred embodiment, the sump temperature should not exceed a value of 160°C.

          Although the method of the invention is not limited to producing a product of any particular content of short and long chain alkyl sulfoquinovosides, in one  
25           embodiment, the product obtained has a high content of the long-chain alkyl sulfoquinovoside and low content of the butyl sulfoquinovoside, alkyl monosulfoquinovosides, and also alkyl poly(oligo) sulfoquinovosides.

          Alkyl sulfoquinovosides are anionic surface-active agents that are suitable for use as industrial surfactants for the manufacture of detergents and cleaning  
30           preparations. Biermann *et al.*, US Pat. No. 5,374, 716, teaches a process for the production of surface-active alkyl glycosides. Miyano, M. & Benson, A.A., "The



Plant Sulfolipid VII. Synthesis of 6-sulfo- $\alpha$ -D-quinovopyranosyl-(1 $\rightarrow$ 1')-glycerol and Radiochemical Synthesis of Sulfolipids," *J. Am. Chem. Soc.*, 84: 59-62 (1962) teaches the preparation of 6-sulfo-D-quinovose from 1,2-isopropylidene-6-O-osyl-D-glucofuranose by sulfate replacement, its subsequent conversion to an allyl  $\alpha$ -glycoside, and its oxidation by permanganate to form sulfoquinovosyl glycerol. Roy, A.B. & Hewlins, J.E., "Sulfoquinovose and its aldonic acid: their preparation and oxidation to 2-sulfoacetaldehyde by periodate," *Carbohydrate Res.*, 302: 113-17 (1997) teaches the preparation of 2-sulfoacetaldehyde by the oxidation of sulfoquinovose, or its aldonic acid, with periodate.

## EXPERIMENTAL

### Example 1

In this example, a means for the production of UDP-SQ from a reaction mixture comprising UDP-glucose, *Arabidopsis thaliana* recombinant SQD1 enzyme protein, and sulfite is described. In one embodiment, the UDP-SQ production reaction is carried out at 37°C in a buffer containing 10  $\mu$ g purified SQD1 protein, 100  $\mu$ M Na<sub>2</sub>SO<sub>3</sub>, 2.2 mM UDP-glucose [<sup>14</sup>C(U)-glucose](69 Bq/nmol) and 50 mM Tris (pH 7.5) in a total volume of 100  $\mu$ l for 40 minutes. The reaction mixture is then heat denatured 5 minutes at 95°C to inactivate the recombinant enzyme, centrifuged at 10,000 X g for 5 minutes, and analyzed by high performance liquid chromatography (HPLC) (Waters Corp., Milford, MA) employing a Beckman (Fullerton, CA) Ultrasphere ODS column (4.6 mm X 25 cm, particle size 5  $\mu$ M) kept constantly at 42°C. Substrates and products were separated by applying a linear gradient of 30 mM KH<sub>2</sub>PO<sub>4</sub>, 2mM tetrabutylammonium hydroxide (Fisher Scientific, Fair Lawn, NJ), adjusted to pH 6.0 with KOH, to HPLC grade acetonitrile (EM Science, Gibbstown, NJ) with a flow rate of 1 ml per minute over 45 minutes.

Incubation of the SQD1 protein with labeled UDP-glucose as described above resulted in the formation of two compounds (U<sub>1</sub> and U<sub>2</sub>) with unique retention times as compared to UDP-glucose (See Figures 2A & B) as analyzed by HPLC. Filtration of the reaction mixture using Amicon filters (MW cutoff 10,000; Millipore Co., Bedford,

MA) without denaturation revealed that 77% of the compound U<sub>2</sub> (See Figure 2B) was free in solution as compared to 35% of compound U<sub>1</sub>. Adding sulfite to the reaction mixture eliminated compound U<sub>1</sub> completely and further stimulated the formation of compound U<sub>2</sub> (See Figure 2C). Compound U<sub>2</sub> co-chromatographed in the HPLC system described above with [<sup>35</sup>S] UDP-SQ indicating that the compound produced in the reaction mixture was UDP-SQ. (See Figure 2D). Labeled compounds were detected using a β-Ram Model 2 Flow Through Monitor (INUS Systems, Tampa, FL).

### Example 2

a. In this example, a means for the production of *Arabidopsis thaliana* recombinant SQD1 enzyme protein, as used in the method described in Example 1, and encoded by the nucleic acid sequence set forth in SEQ ID NO:5, is described. In order to isolate *A. thaliana* genes encoding enzymes involved in the head group biosynthesis of thylakoid membranes, the dbEST database of expressed sequence tags was searched with the predicted amino acid sequence of the bacterial *sqdB* genes using TBLASTN. Through said search, a partial rice cDNA (EST D46477) was found that encodes a putative protein with high sequence similarity to the bacterial *sqdB* gene products. A 400 base pair *Xho* I-*Eco*RV fragment of the partial rice cDNA was used as a probe to screen 2.4 million plaque-forming units (pfu) of an *A. thaliana* PRL2 cDNA library (a lambda ZipLox-based library available from the *Arabidopsis* Biological Resource Center at Ohio State University, Columbus, OH) by heterologous DNA hybridization. Hybond N+ (Amersham) membranes were used, and hybridization was performed at 53°C in 0.25 M sodium phosphate buffer (pH 7.2) containing 7% (wt/vol) SDS, 1 mM EDTA, and 1% (wt/vol) BSA. After hybridization, the membrane was washed twice for 20 minutes in a 2X SSPE, 0.1% (wt/vol) SDS solution at 53°C.

Several cDNA clones were isolated, including one with an insert of 1,799 base-pairs, which was sequenced (GenBank accession No. AF022082). The open reading frame (ORF) beginning at nucleotide 170 encodes a putative protein with a calculated

molecular mass of 53.1 kDa. An amino acid comparison analysis of the *sqdB* gene of *Synechococcus* sp. PCC7942 and the deduced amino acid sequence of the *A. thaliana* cDNA revealed a sequence identity of 42%. The corresponding locus of *A. thaliana* was designated SQD1 and the plasmid containing the cDNA with the 1,799 bp insert was designated pSQD1. At the amino acid level, the partial rice cDNA sequence was 86% identical to the SQD1 sequence of *A. thaliana*.

To produce recombinant SQD1 protein in *Escherichia coli*, a 1,199 base-pair fragment of pSQD1 (nucleotide numbers 425-1603 of GenBank accession no. AF022082) was cloned into the His-tag expression vector, pQE-30 (QIAGEN, Inc., Valencia, CA) using a PCR-based strategy. For this purpose, a forward primer having the nucleotide sequence 5'-AAA GGA TCC CGT GTT ATG GTC ATT GG-3' (SEQ ID NO:10), and a reverse primer having the nucleotide sequence 5'-GTC GGA TCC TTA TGT GGT CAT GGA CT-3' (SEQ ID NO:11) were used such that a *Bam*HI site was provided for cloning into pQE-30, and that the N-terminal 85 amino acids containing the presumed signal peptide were removed. The resulting plasmid construct, pSQD1-TP, allowed the expression of the recombinant SQD1 protein in *E. coli* and the purification of the protein due to the selective binding of the six N-terminal histidine residues of the plasmid construct to Ni-NTA agarose following the manufacturer's instructions. (QIAGEN, Inc., Valencia, CA).

The recombinant protein was eluted with 200 mM imidazole, which was subsequently removed by use of a Millipore Ultrafree 4 concentrator (Millipore, Inc., Bedford, MA). The protein was stored in 20% glycerol, 300 mM NaCl, and 25 mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.5) at -20°C. The SQD1 protein was estimated to be approximately 95% pure by SDS-PAGE gel analysis (See Figure 4).

b. An enzyme assay was developed to measure the conversion of UDP-glucose to UDP-SQ as predicted for SQD1 activity. Basic activity assays were carried out at 37 °C in a buffer containing 10 µg purified SQD1 protein, 100 µM Na<sub>2</sub>SO<sub>3</sub>, 500 µM UDP-glucose [<sup>14</sup>C(U)-glucose](89 Bq/nmol) and 50 mM Tris (pH 7.5) in a total

volume of 100 µl for 40 minutes. An further alternative assay, the coupled adenosine 5'-phosphosulfate (APS)(Sigma, St. Louis, MO)/SQD1 assay, contained 50 mM Tris (pH 8.5), 10 mM dithiothreitol (DTT), 25 µM [<sup>35</sup>S]APS (500 Bq/nmol), 250 mM Na<sub>2</sub>SO<sub>4</sub>, 1 mM EDTA, 500 µM UDP-glucose, 66 µg purified SQD1 protein, and 12 µg APR1 from *A. thaliana*. (See Figure 8). In both assays, the reaction was incubated at 30°C for 10 minutes. The samples were heat denatured for 5 minutes at 95°C, centrifuged at 10,000 X g for 5 minutes, and analyzed by HPLC (Waters Corp., Milford, MA) employing a Beckman (Fullerton, CA) Ultrasphere ODS column (4.6 mm X 25 cm, particle size 5 µM) kept constantly at 42°C. Substrates and products were separated and analyzed by HPLC as described above in Example 1.

### Example 3

In this example, a means for the simplified transformation of *Arabidopsis* is described herein and follows the methods of S. Clough and A. Bent, "Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*," *Plant J.*, 16:735-43 (1998).

a. In this example, a *Agrobacterium tumefaciens* strain carrying the gene of interest, *SQD1*, on a binary vector is prepared as follows. The entire *SQD1* coding sequence (See SEQ ID NO: 6), including transit peptide, but excluding DNA 5 prime of gene, is cloned into pBluescript II (Stratagene, La Jolla, CA) using a PCR-based strategy. For this purpose, said *SQD1* sequence was amplified by PCR using a forward primer having the nucleotide sequence 5'-CTA GGT ACC AAA TGG CGC ATC TAC TT-3' (SEQ ID NO: 20), and a reverse primer having the nucleotide sequence 5'-GTC GGA TCC TTA TGT GGT CAT GGA CT-3' (SEQ ID NO: 11). The primers were constructed such that *Kpn* I and *Bam*HI sites were provided for cloning the *SQD1* cDNA fragment into pBluescript II.

The *SQD1* cDNA fragment is then excised from pBluescript II using the above restriction endonucleases, and subcloned into the corresponding restriction sites on the binary vector, pBINAR-Hyg. This vector is derived from pBIB-Hyg (Becker, D.,

*Nucleic Acids Res.* 18: 203 (1990)) by insertion of the *Hind* III-*Eco* RI fragment from the central portion of pA7 (von Schaeven, A., Ph.D. thesis, Freie Universität Berlin (1989)). This construct is introduced into *Agrobacterium tumefaciens* strain C58C1 and used to transform *Arabidopsis thaliana* Col-2 plants as described below.

5           b.        *Arabidopsis* plants are grown under long days in pots in soil covered with bridal veil, window screen or cheesecloth, until they are flowering. First bolts are clipped to encourage proliferation of many secondary bolts, causing the plants to be ready roughly 4-6 days after clipping. Optimal plants have many immature flower clusters and not many fertilized siliques, although a range of plant stages can be  
10       successfully transformed.

          The *Agrobacterium tumefaciens* strain carrying the gene of interest on a binary vector is grown in a large liquid culture at 28°C in LB (10 g tryptone, 5 g yeast extract, and 5 g NaCl per liter of water) with 25 µg/ml hygromycin B (Calbiochem) to select for the binary plasmid. The *Agrobacterium* culture is pelleted by centrifugation  
15       at 5500 X *g* for 20 minutes, and resuspended to OD<sub>600</sub> = 0.8 in a sterile 5% Sucrose solution.

          Before the above-ground parts of an *Arabidopsis* plant are dipped in the resuspended *Agrobacterium*/Sucrose solution, Silwet L-77 (OSi Specialties, Inc., Danbury, CT) is added to a concentration of 0.05% (500 µl/L) and mixed well. The  
20       above-ground parts of an *Arabidopsis* plant are dipped in the *Agrobacterium* solution for 2 to 3 seconds, with gentle agitation. The dipped plants are placed under a dome or cover for 16 to 24 hours to maintain high humidity. The dipped plants are not exposed to excessive sunlight as the air under dome can get hot.

          The plants are grown for a further 3-5 weeks and watered normally, tying up  
25       loose bolts with wax paper, tape, stakes, twist-ties, or other means. Watering is halted as the seeds of the plant become mature. Once mature, the dry seeds are harvested by the gentle pulling of grouped inflorescences (*i.e.* flower clusters) through fingers over

a clean piece of paper. The majority of the stem and pod material is removed from the paper and the seeds are stored under dessication at 4°C.

Successful transformants capable of expressing a recombinant *A. thaliana* peptide are selected by using an antibiotic or herbicide selectable marker. In this example, 2000 harvested seeds (resuspended in 4 ml 0.1% agarose) are vapor-phase sterilized and plated on selection plates with 50 µg/ml hygromycin B, cold treated for 2 days, and then grown under continuous light (50-100 µEinstein) for 7-10 days. The selection plates of the example are further comprised of 0.5X Murashige-Skoog medium (Sigma Chem. Cat.# M-5519) and 0.8% tissue culture Agar (Sigma Chem. Cat.# A-1296). Successful transformants are identified as hygromycin-resistant seedlings that produce green leaves and with well-established roots within the selective medium.

A sample of successful transformants are grown to maturity by transplantation into heavily moistened potting soil. Leaves from the transformants are removed and subjected to DNA extraction to isolate the genomic DNA of the plant. The extracted genomic DNA is subsequently subjected to restriction endonuclease digestion and Southern Blotting to confirm the incorporation of the gene of interest into the plant's genome.

#### Example 4

In this example, a means for the expression of a peptide, SQDX (SEQ ID NO:1), as contemplated in the example above, is described. The entire insert of the plasmid pSYB carrying the *sqdB* gene of *Synechococcus* was sequenced (GenBank Accession No. AF155063) leading to the identification of a new ORF (open reading frame) directly 3' of *sqdB*. The plasmid pSYB is derived from the plasmid pBlueScript II-SK+ (Stratagene, LaJolla, CA Cat.# 212205) and contains the entire sequence of the *sqdB* gene cDNA (SEQ ID NO: 8) cloned into the plasmid's *KpnI* and *BamHI* sites. This ORF encodes a putative protein of 377 amino acids with no sequence similarity to any of the described *sqd* gene products of *R. sphaeroides*. Unlike the preceeding *sqdB* ORF which starts with GTG, the second ORF begins with

ATG 15 bp from the 3' end of the *sqdB* gene. This ORF was designated *sqdX*. Analysis of the deduced amino acid sequence of *sqdX* (Figure 17: SEQ ID NO: 2) employing Pfam (Protein families database of alignments) revealed a glycosyltransferase group I domain between the residues 228 and 347.

To confirm that the *sqdX* gene in the cyanobacteria *Synechococcus* encodes functionally homologous proteins, the *sqdX* open reading frame of *Synechococcus* was inserted behind the *tac* promoter in the mobilizable broad host range plasmid pRL59EH (Black *et al.*, "Analysis of a Het- mutation in *Anabaena* sp. PCC7120 implicates a secondary metabolite in the regulation of heterocyst spacing," *J. Bacteriol.*, 174: 2352-2360 (1994)), and transferred the constructs by conjugation into *Synechococcus* mutant 7942 $\Delta$ *sqdX* as described in Wolk *et al.*, "Construction of shuttle vectors capable of conjugative transfer from *Escherichia coli* to nitrogen-fixing filamentous cyanobacteria," *Proc. Natl. Acad. Sci. USA*, 81: 1561-1565 (1984). Sequences 5' of the presumed ATG up to the first in-frame stop codon (position 2385912-2387168 of the genome sequence) were included. The *sqdX* gene of *Synechococcus* was PCR-cloned from the plasmid pSYB using the primers 5'-AAG GAT CCT GCG CTA AAG TCG CAC TC-3' (SEQ ID NO: 21) and 5'-ATA AGC TTC GAG CTC AGG CCG CT-3' (SEQ ID NO: 13) into the *Hind* III/*Bam* H I sites of pRL59EH. An  $\Omega$  cassette from the plasmid pHP45 $\Omega$  (as described in Prentki, P. and Krisch, H.M., "In vitro insertional mutagenesis with a selectable DNA fragment," *Gene*, 29: 303-313 (1984)) conferring spectinomycin and streptomycin resistance was inserted into the *Hind* III sites of these plasmids to provide a suitable selectable marker. The resulting plasmid containing *sqdX* of *Synechococcus* was designated pSQDX7942. Exconjugants were selected on BG11 medium containing 25  $\mu$ g/ml kanamycin, 10  $\mu$ g/ml spectinomycin, and 1  $\mu$ g/ml streptomycin and were analyzed by DNA/DNA hybridization to confirm the presence of the proper plasmid construct. The insertion of the *sqdX* construct restored the sulfolipid biosynthetic activity in the *Synechococcus* mutant 7942 $\Delta$ *sqdX* as shown by TLC lipid analysis. Based on the

observed genetic complementation, it is concluded that the cyanobacterial *sqdX* gene encodes a protein involved in sulfolipid biosynthesis.

#### Example 5

In this example, a means for the production of *sqdX* gene homologs of *Arabidopsis thaliana* comprising the group consisting of the gene product ATSQDX-1 encoded by the nucleic acid sequence set forth in SEQ ID NO: 3, the gene product ATSQDX-2 encoded by the nucleic acid sequence set forth in SEQ ID NO: 4, or the gene product ATSQDX-3 encoded by the nucleic acid sequence set forth in SEQ ID NO: 5, is described. A BLAST comparison of the cyanobacterial *sqdX* gene to genomic sequence of *Arabidopsis thaliana* revealed several potential homologs. In one example, *AtSQDX-1*, a homolog having 37% amino acid identity with the cyanobacterial *sqdX* gene is contemplated. In another example, *AtSQDX-2*, a homolog having 29% amino acid identity with the cyanobacterial *sqdX* gene is contemplated. In a further example, *AtSQDX-3*, a homolog having 32% amino acid identity with the cyanobacterial *sqdX* gene is contemplated.

#### Example 6

In this example, a means for the subsequent modification of UDP-SQ to produce an alkyl sulfoquinovoside is described. The synthesis of alkyl sulfoquinovosides starts from the hydrolytic cleavage of the UDP-SQ. Sulfoquinovose is then refluxed with the acidic catalyst, para toluenesulfonic acid, in the presence of the short chain alcohol, butanol, to form a short chain butyl sulfoquinovoside. The reflux temperature is 118°C. With the formation of the lower boiling butanol/water mixture, a vapor temperature of 95 to 110°C is established. The acetalization with the butanol is carried out under light vacuum, i.e. under a pressure of 800 to 950 mbar. An azeotropic amount of butanol is removed with the water in the distillation process.

The butyl sulfoquinovoside is subsequently treated under vacuum with the long-chain alcohol, n-hexadecyl alcohol, in the presence of the acidic catalyst to form a long-chain sulfoquinovoside. In order to obtain a low content of butyl






in *Molecular Biology*, John Wiley & Sons, NY (1992)). After expression, the protein is isolated and purified. The protein may then be used for the generation of antibodies (see, generally, Howard and Bethell, *e.g.*, *Basic Methods in Antibody Production and Characterization*, CRC Press, (2000)).


5 Alternatively, preparative reagents are generated to isolate the specific target by conjugating antibodies generated from expression of fragments of the genomic sequences known to contain the desired sequence. The antibodies are generated by methods known to those in the art (*See, generally*, Howard and Bethell, *e.g.*, *Basic Methods in Antibody Production and Characterization*, CRC Press, (2000)) to generate  
10 anti-AtSQDX antibodies. The antibodies are conjugated to agarose beads. Furthermore, a parallel conjugate of agarose beads to a control immune globulin is accomplished. Ultracentrifuged cell lysates from the desired cell line are exposed to the control non-immune I conjugated beads to remove non-specifically binding proteins. The unbound lysate is recovered and is then exposed to the anti-AtSQDX  
15 antibody conjugated agarose beads for a direct affinity purification. The anti-AtSQDX antibody/AtSQDX complex is washed with 2.5 M KCL to remove non-specifically bound materials and the AtSQDX is then eluted from the agarose beads with 0.1 M glycine HCL in the presence of 0.5 M NaCl. The eluted material from the column is neutralized with 1 M Tris pH 8.0, dialyzed extensively to reduce the salt concentration  
20 to 150 mM and then reconcentrated. The reconcentrated material is placed on SDS-PAGE under non-reducing conditions for a final purification based on molecular size. This material is transferred to a membrane for electrospray tandem mass spectroscopic analysis of the amino acid sequence. This later sequence is used to generate oligonucleotide probes for the cloning of the gene encoding AtSQDX.



Figure 1




**Figure 2**




# Figure 3

## pQE-30, pQE-31, and pQE-32 Vectors


Positions of elements in bases	pQE-30	pQE-31	pQE-32
Vector size (bp)	3462	3464	3463
Start of numbering at <i>Xho</i> I (CTCGAG)	1-6	1-6	1-6
T5 promoter/lac operator element	7-87	7-87	7-87
T5 transcription start	61	61	61
6xHis-tag coding sequence	127-144	127-144	127-144
Multiple cloning site	145-192	147-194	146-193
Lambda <i>t</i> <sub>0</sub> transcriptional termination region	208-302	210-304	209-303
<i>rrnB</i> T1 transcriptional termination region	1064-1162	1066-1164	1065-1163
ColE1 origin of replication	1639	1641	1640
β-lactamase coding sequence	3257-2397	3259-2399	3258-2398




### pQE-30



### pQE-31



### pQE-32



**Figure 4**





Figure 5





# pACYC184

4,244 base pairs  
GenBank Accession # X06403

Figure 6





Figure 7



UDP-Sulfoquinovose

Butyl-Sulfoquinovoside

Alkyl-Sulfoquinovoside

**Figure 8**




Figure 1 consists of two panels, A and B, showing SDS-PAGE gels of the purified DGDG synthase complex. Panel A shows a 12.5% SDS-PAGE gel with three main bands labeled from top to bottom: U<sub>2</sub> Product, UDP-Gal, and UDP. Panel B shows a 12.5% SDS-PAGE gel with five main bands labeled from top to bottom: MGDG, PG, DGDG, SQDG, and PC. Both panels include molecular weight markers on the left side, indicated by numbers 100, 75, 50, and 37.

44

Figure 10

SQDX

1							
2-61	cgcatcgc	tctctttacc	gagacgttcc	tccccaaagt	ggatggcatc	gtcacgcggc	a
62-121	ttcggcacac	ggtcgatcac	ctgcagcgtc	ttggccacac	cgtcatgggt	ttttgccccg	
122-181	acggcgggct	ccgcgagcac	aagggggctc	gagtcctatgg	ggttaaaggc	tttccgctac	
182-241	cgctctatcc	cgagctgaag	ctagcttttc	cgttgccgaa	agtgggaaaa	gccttgagac	
242-301	ggttcgggcc	cgacctgac	cacgtggtca	atccggctgt	gttgggggtg	ggcggcatct	
302-361	actatgccaa	ggcgctaaat	gtgccactcg	tggcgtccta	tcacacccat	ttgccgaaat	
362-421	accttgagca	ttacgggctg	ggggtcttgg	aggggggtgct	ctgggaattg	ctgaagctgg	
422-481	cgcataacca	agcagcgatc	aacctctgta	cttcaaccgc	gatgggtgcag	gagctgacag	
482-541	atcacggcat	tgagcactgt	tgctcttggc	agcgaggagt	ggataccgag	acctttcggc	
542-601	cagacttggc	tactgctgcg	atgcgcgac	gcctcagtgg	cggtaagccc	actgcgccct	
602-661	tggttgcctc	cgtcggacgc	ctctcagccg	agaagcaaat	cgatcgctg	cgacccattt	
662-721	tggatgccaa	tcctgaggct	tgcttggcct	tggtcggcga	tggcccgc	cgggcccgaac	
722-781	tagagcaatt	gtttgctggc	accagacgc	agttcattgg	ctatctgcat	ggggaacagc	
782-841	taggggcggc	ctacgcttct	gctgacgctt	ttgtctttcc	ctcccggacc	gaaaccctcg	
842-901	gtctagtctt	gctggaagcc	atggcagcgg	gttggtccgg	cgtggcggcc	aattccgggtg	
902-961	gcattcccga	tattgtcagc	gacggcatta	atgggtttcct	gttcgatcct	gaggatgaac	
962-1021	aaggggcgat	cgctgcgatt	cagcgcttgt	tggctaacc	tgcagagcgc	gagattctac	
1022-1081	gccaagcggc	tcgtcaagaa	gccgaacgct	ggagctggaa	cgcagccacg	cgccaactcc	
1082-1134	aggactacta	ctgcgaggtg	ttggcagatg	gttgcttacc	cttagcggcc	tga	

05709020.1.10300

### Figure 11

AtSODX-1

1-57	ctacacg	ttaccttccg	gtactggaaa	cagtcgttta	atcaaccaat	tgattggtcc
58-117	caaaacatga	actttctttt	tcctccagaa	ccaaatcgct	gcactgtact	gttcatttgcg
118-177	tatctttgtc	gttgctgctc	tccaatcata	tttctcgggc	tcttctcttg	ccgcttttcc
178-237	aatgatctct	cttgtttcac	ggtcgtgcag	taaagtcttc	agttttgtca	cgcaatcttc
238-297	aacatctcca	gggttgaaca	aaaatccggt	ttttccctcc	tgatgaaaac	atcagaatca
298-357	gaaaaccaca	agctcaatat	aggttgaccc	ataagaacaa	tcaatgcaag	atcattttgt
358-417	gtaccagtct	atgattgaat	aaagtttcag	ttcggttaca	gctcgcttat	aagaaaattg
418-477	gcagaaattg	ttttttcaac	catttcgggt	cgggtgatat	gctcatcaat	atgggttggc
478-537	agttaattgt	aattcagata	attcactgac	ctgatcttca	gggattatat	cagggattcc
538-597	accggcacgg	gccgcgacga	caggaagtcc	tgaagacatt	gcttcaagaa	ccacaaggcc
598-657	aagtgtctcc	gactctgatg	gcatacacia	cacatctcca	cttgcgtaag	cttgtgagag
658-717	ttcatcgctc	tgtaacggtc	cagtgaaaac	cgctggcatt	ccggtaaaaa	acttctcaag
718-777	atcctctctt	taagaaaaag	aaacagataa	acaaaattac	aatggtgttg	actagaaatc
778-837	ttcagataac	aatatggcca	atctttaaca	aaactagtag	ttgtatgggc	catctccaat
838-897	gaaagcaatc	cgagcttcag	gtaatttgtc	cattacactg	cacacaaatt	tctcaatatc
898-957	aaaattcgat	acaccactta	aaagaagtga	gtccagttta	tacaaaaatt	taacctcttt
958-1017	aaaagctcca	aactcttttc	tacgccaatg	cgacctacat	gaatcactag	tggcttttct
1018-1077	ggttcgccat	tactgttaat	tacaaaatat	taaacatcaa	gattagcgtg	gaaagtatca
1078-1137	ttgttttttaa	tgcatataaa	agaaacgtat	attctattct	tgcttcagtc	ttatacgcat
1138-1197	ttcttgagaa	cggaaacggg	gattgaagct	ttctgaatcg	acaccttat	tccaaagtgc
1198-1257	aagttgatta	gcttcaatat	atgaaaaaaa	gaagaagaaa	atgtaagttt	tgaacaatca
1258-1317	tagagcttgt	acaaaaatgt	aatgtatgat	cttcttcac	ctgcagttgc	accagctgct
1318-1377	ataagatctt	ttccaatggc	agcagaagga	actaatgtaa	gatcagccgc	tctgtgaagg
1378-1437	aaccttgata	aaagcatatt	caagtttagt	ttcatattat	acatcacaat	aaaccagaaa
1438-1497	aagaagaagg	aaattttgac	atttgaaaag	cgggttttac	atacttatta	tagaccacat
1498-1557	tggtttttacc	aaccaactaa	aagtgtatct	tgggatgtat	ctgcacatca	acacaagctt
1558-1617	aatcttagac	aaaaattttt	ttataacaac	attgtgaaat	gaggcagaaa	aaggtactta
1618-1677	cacagggacg	tgtgtgtggg	aagacattac	tattggtaca	gatagcattt	ttgctattgc
1678-1737	cagagcacca	aagacctaaa	aatttttagtc	agggaaaaaa	gagagtcaag	ttctggattc
1738-1797	tctccagttc	actggtcttg	ctaattgttt	agtaatgtga	attcttgagg	gatttaccat
1798-1857	aactccggga	gatgaagcgt	gtataatgtc	aggcttaaac	cgtgcaattt	cagagatgat
1858-1917	tcttggaacta	agcgcaagcg	agagtggaa	cttttggtaa	taggacaag	ggaagctaca
1918-1977	gaagagaga	agaaattagc	gatattacca	aatagagaac	atccagttag	taaactaaat
1978-2037	gggtgctacct	tcttgatcca	atgactctgg	ctccataaaa	ctcttcagga	acaccttcac
2038-2097	gtgtcgtcac	gactataaacc	tatgaagcaa	aaaagtattt	aaaaaaaaaa	aaaaaggaac
2098-2157	agttaacact	tgtcaagtaa	ttctaattct	ggaaacagtt	acttatgagc	tgactgaaaa
2158-2217	gatacttaag	ttgaagaatg	agatagtaaa	agaagaaacc	tcgtctccca	tttcacggag
2218-2277	gtatctaatg	aaattctgga	atctgttttt	gtagccggat	acatagctgc	aacaaaatca
2278-2337	aagagagaat	cacttccaat	aataacatga	catataataa	aagcttttgg	tcaatggatc
2338-2397	gggtgattccg	agaatcttgg	gatattcaca	actaaaatct	gacaactttg	actcaaacaa
2398-2457	atcctgaatg	taattggttt	taacgatcta	ctatataatt	tgctaaattg	gtgggtgtagc
2458-2517	aaattcatac	attagcgagt	atctcttcat	aaaataaatg	taacgatcaa	atcgaaagaa
2518-2577	aaaaacatta	caggaaaaag	tcaaccaagg	aaaaaaatga	gtagaatctg	ttttcacaga
2578-2637	gacatttcgt	cgaacacaaa	acaagcaaaa	aaagaacact	gtgaagaaga	cttacgcaaa
2638-2697	gggagaaggc	tcaacaaaga	gagcaattct	cctaggctta	gagagcgact	caggatcgag
2698-2757	aagcggcgca	tcaatctccg	attcgtcatc	ttctctgact	tgagtaatag	tcatatcggt
2758-2817	ggaccagaa	acagcttcct	ttgtaatcac	tccacagaag	cggagcttgc	ttttcttgct
2818-2877	aatgggtaat	ctccggtgac	caaacgaaag	aggggaatga	agaacaaaag	aagaagacct
2878-2937	gggaggagaa	caagaggttg	cagaggaaga	agaacaagtg	ttagtcgtgc	taggaagcaa
2938-2979	atgaggaggt	atagagagat	ttatagaaga	aaqaqtcgtc	at	

# Figure 12

AtSQDX-2

1-50		tcatattttg	aaaaagcact	tcttcttctt	cttcttcttc	ttcttcttct
51-110	tcttcttccc	atcaaaaagt	gctagtggca	agaccgcgca	tattccagta	acaaagttta
111-170	aattcaaaac	tatggaacac	aatttgctcc	atgttttggt	gcctgaatta	aacgaccata
171-230	tcacttggtt	cggccacttc	atctcggata	cgcacaagcg	gttgtcaagg	ttgtaaaaga
231-290	cgattttgta	aggatctaca	ttggcaaagg	gagctttaca	gacgacttga	aaagtttcgg
291-350	tgtgaagatt	gaaaggtaaa	acttttggtt	ctttgcactc	ggtgaaccaa	taaagcgacc
351-410	catctagata	cacaggtgcg	gggtaagcag	caatccgata	aggagcagcg	agagtgcacat
411-470	acctccaagc	attgggtacta	aagtcgaaaa	cttcgcgatgt	agtagcggtt	tttaggccta
471-530	tttctgcaga	gttgtataac	caaacgggct	tgtatgtgcc	cctgaatttg	tctttaccga
531-590	atccaagcat	aaagaatttg	cgtttaagct	tgtagtaacc	atctcgtaag	togatcatga
591-650	gttttttgata	atcgcaaaga	ggaagaggcc	gataccatct	agtggtcggg	ttgaccacat
651-710	aaccggattt	gtcgtgattg	taaagacaaa	cgagaccgtc	acaactacta	tgtgaaacta
711-770	agtacagtac	gttatccttc	tcccaaggag	tagggatctt	gaacgttgat	gatgaaccca
771-830	actgcaatgt	tcttaaagat	tctatggtcg	ggtttataac	atggtgagga	aatacagaca
831-890	ccatcagaac	atctggatct	cctgattgct	gacgatgttt	caactaaaat	tgtatacatt
891-950	tttattttat	taaaaagata	aatatcattt	tgttggaatt	ggaaaacaaa	atctaaaaga
951-1010	taacttttaga	tatgtaaggg	tcgcatatgt	atttgattac	attgctatga	gtatatgact
1011-1070	cacttttcca	ataatgaaaa	aaataaagat	gttgagtctt	actaattaag	ggtaaataca
1071-1130	aaattttctga	tcattaatac	aaagaaaaag	cctctcataa	gccaaagcca	ttttactcgc
1131-1190	cgtaaaccatc	tccgcagcat	actctttaca	cttccttcct	ctctccgcca	atctctccgc
1191-1250	tccttcagcc	acagcaacct	ccataaccgc	cgtcaaagct	tccacatttg	gcgcaaacat
1251-1310	aaacccaaac	tcatcattca	ccactatagt	cctctttatg	cttgcgtagc	tagatgccat
1311-1370	cactggttta	ccactcaaca	tcgcttccat	taacgttaaa	tcaagacctt	gtggtctaag
1371-1430	cgttggattc	acgaacaaat	cgatcccgtt	gtagaaaccc	ttgagctcat	ttgggtttag
1431-1490	agatcccaaa	atggaaactt	tttcccctaa	ttccttgtaa	cgttgctccc	atggccctga
1491-1550	tccagctact	acgaggtaaa	catttgaaata	cgtttggtat	attttcgcga	aagcttcgaa
1551-1610	gagcaatgga	tgtcctttgt	ctttgactaa	tctcccagca	gctcctaaaa	caatcgctga
1611-1670	tgagttttct	ggtaacccta	attttgacct	aaacagagta	cgtagcttct	tgtctgatgt
1671-1730	gaatccgttc	tcgtcgactc	cattgaggat	cacatgaacc	cttttctcag	ggatttggtg
1731-1790	aacgtctctt	agcattttctc	cgcagctatc	gctgatagcg	atgtggtagg	cgtagttgtg
1791-1850	gaagaatctg	atttcgtcca	gtatcttggg	aagcacggca	ccgtatagac	ttgcattgaa
1851-1910	gccttgatgt	cttggttcgt	ctgggtttacg	gatcaggtct	tggtaaatac	ttgattgtaa
1911-1970	gctctctaac	gcaatgccgt	gccaggatac	agcgagggtt	ggaacctccc	gggcgatcca
1971-2030	gtgaggtaaa	gcaacacttt	cagagtgaac	cgcacgaaa	ggttctttct	tgttttcttc
2031-2090	ttggtaaagc	tcccatgcct	tgttgtagccg	ccattttccg	ggctccgctg	cgccgtgaga
2091-2150	atggattata	gggtaaatga	tttggttcgga	aacagggggg	attttggttg	tttcagggga
2151-2210	ttggtctaag	ggagaggtga	aaacgtggac	acggtgtcca	cggcgagcta	aggcggtgta
2211-2270	tagagtgaag	gcgtggcggt	ccatgccgcc	aggattggga	cccgtgggcc	atttccggga
2271-2330	gaaaacagcg	agtttttagtg	ttttggggagg	agggttggtt	aaggagaaat	caaggcggtt
2331-2390	ccaggcaaat	tgggcggttt	ggaggtcgcc	agaccacggt	ggctgggttg	tgtcagagga
2391-2450	ggaggaggag	actgcagcgg	tggaggagga	gcaagtggag	gtgctggagga	gaaagagagc
2451-2510	cggtatggta	aagaggacgg	tgaagaagag	gaaagtacaa	aggctaaaat	gagaattagg
2511-2537	tttcttgagt	ttggtttggtg	aagccat			

# Figure 13

AtSQDX-3

1-7						ttaaggt
8-67	ctcatgcatt	tgaccagaac	atcgacgaat	cgcttgtaca	tatgttgctt	catgtacatt
68-127	ttctcaacca	ttttgcgtcc	ttcactcccc	aatcgtagcc	tctcgtctgg	attcctaagt
128-187	agatacaaga	gattatgagc	taattctttg	ttaccggatc	tccccattga	gtgaagtaga
188-247	ccagtcattg	tgtgttgaac	catctctttg	gttcctcctg	catctgttcc	caccactgca
248-307	agtccataag	ccattgcttc	gattgtcact	ctaccaaag	tttcaccaac	tccctgtaaa
308-367	aaatcaaggt	tcaaaataaa	taaaccgaac	aaaatcgaaa	tggtcaaacc	gaaatcaaaa
368-427	cctacatttt	gaacaaaaac	aaaactgaaa	tgttttatcc	aaaaccaaac	tgaaatcaat
428-487	cacaaattgg	tttatttgat	tcagtaattc	aagttcctat	aaaactgata	aaactaaacc
488-547	aaaacctaac	tgtaaactac	tttctagtgg	aaaactgcat	atgcatcaaa	taatgttttt
548-607	gaggtgtgag	gaggattacc	tgggagtttg	taacgtagac	atctgctgca	gagtataatg
608-667	aagcaacacg	ggttgttgca	ggagtccaca	ttaccgactt	agataagttt	ccgctgtttg
668-727	acaagaagct	taacatctct	ttaacgtatc	caactttgtt	gctctttgaa	cccacggatc
728-787	ctaaaagaac	tttaagttct	tgcttctctc	ttcttaagcc	attgtcaagt	gtgagagaaa
788-847	cacttttcat	ctgtcgtgat	gaacctctta	aacgatgttt	gctggagaga	ctaacctttt
848-907	ctttccta	gatccccctg	tgattccttt	gagattcctt	tcctctctca	gaaagagcca
908-967	aggcaataga	ttcaaggaga	agaagttgtc	cctttgttgg	gtttatgctg	ctaagagaca
968-1027	tcacaagcat	atctgaatct	gttattccta	actctgttct	cactgattcg	cgtaatat
1028-1087	gtctcttcac	cctcattttc	tctggtgaaa	gtgttggtgt	gttgagtga	gaaggaatcc
1088-1147	ccgctacaaa	agctaactca	tcattaacag	atagtggaa	aatcactgg	tgtgatctaa
1148-1207	gcttttatat	ctcctcctcg	caccatgtta	accattgtct	gctctgtgat	tcagataaga
1208-1267	aaatcagcat	tttcactcgg	tcaagaactg	gtttcgctcg	atcaaagtat	tctcgtcgat
1268-1327	tctccattat	ccaccaagct	atgtgacttc	caccagctgg	atgatgattc	atgtattgat
1328-1387	ctgcgagaaa	aggaaaaaaa	aaaagtaatt	aagatatgct	ttcctctgat	tgtaaattag
1388-1447	tagtctaaaa	actgcataat	gaattcttac	ctatccatga	ggtacacact	gctgatcctg
1448-1507	cgatgatcaa	atctgctttc	atggcagtct	tgaagctgag	ttctcctttg	tcttcaacaa
1508-1567	ctttgatcct	tctcctactg	agctcttgca	tcaatccacc	tctcctgcta	agaactactg
1568-1627	cagagactgt	tgcaccacag	ctcaaaagct	ctgaagccag	ctccatcata	gaaattggag
1628-1687	caccagtcatt	tgatagctcg	tggaaaagca	ggacgaatct	ccttgaccaa	acaaggcggt
1688-1747	taaaatctga	tttcctgtcg	caagtcccag	atcttctatg	cgggctccat	tcaagaactt
1748-1807	tatcctctag	tgatccaaag	ggaccaagaa	gcttaccata	agtagcattt	gtcaatggaa
1808-1867	gttggtggatc	ttgctcatca	tccaaatcct	tagtctcaag	tacctcttta	atcaccttct
1868-1927	gcttaacacg	gatcttacta	cgagaagtcc	ggacagtttt	tctcgtcttc	tgcttggaac
1928-1987	tcaagctccg	tcgagaaacc	ccatcatctt	tcttgatcag	actaacatcc	gtcctcttat
1988-2047	tcgaaccagc	atcatcttta	ccagtaatat	taaccaaagc	ctcagaatcc	tcattagcaa
2048-2107	caacatccag	tcctttaatc	ttctccatat	ataactcatc	ctttctcggt	ctgcctccaa
2108-2167	accgtaaaaa	ctcaactttg	ctttcattat	catgtgcca	cctagactga	acataaaaac
2168-2227	caagatacgt	caaagcgta	atcaaaagca	gccaataaac	caaccggcta	ctacgaaacc
2228-2287	actgaaacgc	tcctcctcca	ccatgacctc	tacgcggagt	cctaccagaa	tacactctag
2288-2347	gtgtaccctt	tggagtagac	ctccttgaca	gtgaagactt	aacacttgct	tgtctcagcg
2348-2369	gcgataaccg	aatctcctcc	at			



Figure 14

SQD1

1 gtcgacccac gcggtccgctc atctctcatc gttccgggag aagagaagag agacccatcc  
61 ctcaacttcaa agttcaaagt ctccaaggat cttctccaac tctctctaaa caagattcca  
121 aatttttcaaa ggtgaatttg tttgatagaa tcaagaacaa accttttaaaa tggcgcatct  
181 acttttcagct tcatgccctt cagttatctc acttagcagc agcagcagca agaattcagt  
241 taagccggtt gtttcagggc agaccttctt caatgctcag cttcttttcaa gatcttctct  
301 caaaggactt ctcttccaag agaagaaacc gagaaaaagc tgcgtttttca gagcaactgc  
361 tgtacctata acccaacaag caccaccgga aacatctacc aataactcat cctctaaacc  
421 aaagcgtggt atgggtcattg gtggagatgg ttattgcggt tgggctactg ctctccactt  
481 gtccaagaag aattacgaag tttgcattgt tgacaacctt gtaagacgct ttttcgacca  
541 ccagcttgga cttgagtcac tgactcctat tgcctccatt catgaccgaa tcagccgatg  
601 gaaggctttg acagggaaat caattgagtt gtacgttggg gatattctgtg atttcgaatt  
661 cttagctgag tctttcaagt cttttgagcc ggattcagtt gtccactttg gggaacagag  
721 atccgctcct tactcgatga ttgaccgggc cagagcagtt tatacacagc acaacaatgt  
781 gattgggact ctcaacgttc tctttgctat aaaagagttt ggagaggagt gtcactttgt  
841 aaaacttggt acgatgggtg agtatggaac tccaaatatt gacatcgagg aaggttatat  
901 aaccataacc cacaacggta gaactgacac tttgccatac cccaagcaag ctagctcctt  
961 ttatcatctt agcaaagtgc atgattcgca caacattgct tttacttgca aggcttgggg  
1021 tattagagcc actgatctca accaaggagt tgtttatgga gtgaagactg atgagacaga  
1081 gatgcatgag gaactccgta accgactgga ttacgatgct gtgtttggta cagcacttaa  
1141 ccggttctgt gtgcaagctg ctgttgggtc cccacttaca gtttatggta aaggtggtca  
1201 gacgagaggc tacctcgata taagagacac ggttcaatgt gttgagatcg ctatagcaaa  
1261 cccggcaaaa gctggtgagt tccgggtctt caaccaattt acagaacagt tttcagtcaa  
1321 tgaactggct tcaactcgta ctaaagcggg ttcaaagctt gggctagacg tgaaaaagat  
1381 gacggtgcct aacccgagag tggaggcaga agaacattac tacaacgcaa agcacactaa  
1441 gctgatggaa cttggacttg agcctcacta tctatctgac tcaacttctg attcgttgct  
1501 caactttgct gttcagttta aagatcgtgt ggacacgaaa caaatcatgc ctagtgtttc  
1561 ctggaagaag attggcgtca agactaagtc catgaccaca caaagtgcag accaatatta  
1621 cacataagga gagattatga aagagatgat gtgttgtttg gtatcttcaa acttcatttc  
1681 tgcaaaagac ttgctaggct taagaggttt tgtccatatt acattgtgca ggttctttta  
1741 tgttagatct taatttcgat gaaaaaaaaa aaaaaaaaaa aaaaaaaaag ggcggccgc

09709090.10900

Figure 15

SQDB

1-25				gtgaa		
26-85	atggtttctg	cgggtggccc	tgcgtctca	atttggctgc	tgcaggtcac	gccgtcacca
86-145	ttgttgacaa	cctcgttcgc	cgcaagacag	acgtggaatt	gggggttcag	tccctcactc
146-205	cgatcgcgac	gattgaacgc	cggttgaagg	catggcaaga	aacgggagg	cagccgatta
206-265	gctttgtcaa	tctcgactta	gcggctgatt	acgatcgct	ctgtgcacta	ctgctagaaa
266-325	cgcagccgga	tgcgatcggt	cattttgccg	aacagcgcg	cgccccctat	tcaatgaaga
326-385	gtgcatggca	taagcgcttc	acgggtcaata	acaacgtcaa	cgccaccac	aatctgctct
386-445	gcgcctgtgt	ggatgtcggt	ctcaagtccc	acattgtcca	cttgggcacc	atgggcgtct
446-505	atggatacgg	tagccatcgc	ggggctacga	ttcctgaagg	ctacttagaa	gtggaagtcg
506-565	tccagcggga	tggccaacgc	tttgaagaga	agattcttca	ccccgttgat	ccgggtagcg
566-625	tctatcacat	gaccaagacg	ctggatcaat	tgttgttcta	ctactacaac	aagaacgaca
626-685	acatccaagt	caccgacctt	caccagggtg	ttgtctgggg	cacgaacacc	gatcactgta
686-745	atctccaccc	ggatctgacc	aatcggttcg	actacgacgg	tgattacggc	acagtcttga
746-805	accgcttctt	gatgcaggcg	gcgatcggct	atcccttgac	tgtgcatggc	gttgggtggcc
806-865	aaacccgagc	cttcatccac	attcgcgact	cagtgcgctg	cgtccaactg	gcgatcgaaa
866-925	atccgccagc	agccaatgaa	aaagtccgca	tctttaacca	gatgacggaa	acctaccaag
926-985	tcaaggattt	ggcagagaaa	gtggcggtat	tgaccgggtg	tgaaatcgcc	tacctgcca
986-1045	atccacgcaa	ggaagccctt	gagaacgact	tgattgtcga	caaccgctgc	ttgattgatt
1046-1105	taggcctcaa	tccgaccacc	ttggacaatg	gcctgatgag	cgaagtggta	gaaattgcgc
1106-1165	agaagtttgc	cgatcgctgc	gatcgcgcca	aaattccctg	cgtttctgcc	tggaccgta
1166-1209	atcaagctga	agctctcagc	gctcctgaaa	ccgctctgcg	ctaa	

**Figure 16**

MAHLLSASCPSVISLSSSSSKNSVKPFVSGQTFNAQLLSRSSLKGLLFQEKKPRKSC  
VFRATAVPITQQAPPETSTNNSSSKPKRVMVIGGDGYCGWATALHLSKKNYEVCIVDN  
LVRRLFDHQLGLES LTPIASIHDRISRWKALTGKSIELYVGDICDFEFLAESFKSFEP  
DSVVFHGEQRSAPYSMIDRSRAVYTQHNNVIGTLNVLF AIKEFGEECHLVKLGTMG EY  
GTPNIDIEEGYITITHNGRTDTLPYPKQASSFYHLSKVHDSHNI AFTCKAWGIRATDL  
NQG VVYG VKTDETEMHEELRNRLDYDAVFGTALNRF CVQAAVGHPLTVYGKGGQTRGY  
LDIRDTVQCVEIAIANPAKAGEFRVFNQFTEQFSVNELASLVTKAGSKLGLDVKKMTV  
PNPRVEAEEHYNAKHTKLMELGLEPHYLSDSLDSLLNFAVQFKDRVDTKQIMPSVS  
WKKIGVKTKSMTT

008077-02060760

**Figure 17**

MRIALFTETFLPKVDGIVTRLRHTVDHLQRLGHTVMVFCDGGLREHKGARVYGVKGF  
PLPLYPELKLAFPLPKVGKALERFRPDLIHVVNPAVLGLGGIYYAKALNVPLVASYHT  
HLPKYLEHYGLGVLEGVLWELLKLAHNQAAINLCTSTAMVQELTDHGIEHCCLWQRGV  
DTETFRPDLATAAMRDRLSGGKPTAPLLLYVGRLSAEKQIDRLRPILDANPEACLALV  
GDGPHRAELEQLFAGTQTQFIGYLGHEQLGAAYASADAFVFPSTETLGLVLLLEAMAA  
GCPVVAANS GGIPDIVSDGINGFLFDPEDQGAIAAIQRLLANPAEREILRQAARQEA  
ERWSWNAATRQLQDYyceVLADGCLPLAA

bioRxiv preprint doi: <https://doi.org/10.1101/2020.04.01.724441>; this version posted April 1, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

**Figure 18**

MKILVLGGDGFCGWPCALNLAAAGHAVTIVDNLVRRKTDVELGVQSLTPIATIERRLK  
AWQETGGQPI SFVNLDLAADYDRLCALLLETQPD AIVHFAEQRAAPYSMKSAWHKRFT  
VNNNVNATHNLLCACVDVGLKSHIVHLGTMGVYGYGSHRGATIPEGYLEVEVVQRDGQ  
RFEEKILHPVDPGSVYHMTKTLDQLLFYYYNKNNDNIQVTDLHQGI VWGTNTDHCNLHP  
DLTNRFDYDGDYGTVLNRFLMQAAIGYPLTVHGVGGQTRAFIHIRDSVRCVQLAIENP  
PAANEKVRI FNQMTETYQVKDLAEKVAALTGA EIAYLPNPRKEALENDLIVDNRCLID  
LGLNPTTLDNGLMSEVVEIAQKFADRC DRAKI PCVSAWTRNQAEALSAPETALR

003017 02050260

## DECLARATION FOR PATENT APPLICATION

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated next to my name. I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled **Compositions And Methods For The Synthesis And Subsequent Modification Of Uridine-5'-Diphosphosulfoquinovose (UDP-SQ)**, the specification of which is attached hereto. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under § 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of Sole or First Inventor: **Christoph Benning**

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Residence: 674 Wayland, East Lansing, Michigan 48823 Citizenship: Germany  
Post Office Address: 674 Wayland, East Lansing, Michigan 48823

Full Name of Second Joint Inventor: **Sherrie Lea Sanda**

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Residence: 5931 Boise Ile Drive, Apt. 70, Haslett, Michigan 48840 Citizenship: United States Of America  
Post Office Address: 5931 Boise Ile Drive, Apt. 70, Haslett, Michigan 48840

Full Name of Third Joint Inventor: **Bin Yu**

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Residence: 1403-A, Sparten Village, Michigan State. U., E. Lansing, Michigan 48824 Citizenship: China  
Post Office Address: 1403-A, Sparten Village, Michigan State University, East Lansing, Michigan 48824

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Christoph Benning *et al.* Group No.:  
Serial No.: Examiner:  
Filed:  
Entitled: **Compositions And Methods For The Synthesis And Subsequent Modification  
Of Uridine-5'-Diphosphosulfoquinovose (UDP-SQ)**

## POWER OF ATTORNEY BY ASSIGNEE

Assistant Commissioner for Patents  
Washington, D.C. 20231

Michigan State University, as Assignee of record of the entire interest of the above-identified patent application, hereby appoints the members of the firm of MEDLEN & CARROLL, LLP, a firm composed of:

Virginia S. Medlen	(Reg. No. 32,050)	Maha A. Hamdan	(Reg. No. 43,655)
Peter G. Carroll	(Reg. No. 32,837)	J. Mitchell Jones	(Reg. No. 44,174)
Kamrin T. MacKnight	(Reg. No. 38,230)	David J. Wilson	(Reg. No. 45,225)
David A. Casimir	(Reg. No. 42,395)	Jason R. Bond	(Reg. No. 45,439)
Jaen Andrews	(Reg. No. 35,051)		

as its attorneys with full power of substitution to prosecute this application and transact all business in the Patent and Trademark Office in connection therewith.

Please direct all future correspondence and telephone calls regarding this application to:

Peter G. Carroll  
MEDLEN & CARROLL, LLP  
220 Montgomery Street, Suite 2200 Telephone: 415/705-8410  
San Francisco, California 94104 Facsimile: 415/397-8338

I hereby certify that the Assignment document filed with the application or filed subsequent to the filing date of the application, has been reviewed and I hereby certify that, to the best of my knowledge and belief, title is with Michigan State University.

Dated: \_\_\_\_\_

By: \_\_\_\_\_

Name: \_\_\_\_\_

Title: \_\_\_\_\_

Michigan State University  
238 Administration Building  
East Lansing, Michigan 48824-1046

**ASSIGNMENT**

WHEREAS, WE, Christoph Benning, Sherrie Lea Sanda and Bin Yu, hereinafter referred to as "ASSIGNOR", have invented certain new and useful improvements as described and set forth in the below-identified application for United States Letters Patent:

Title of Invention: **Compositions And Methods For The  
Synthesis And Subsequent Modification  
Of Uridine-5'-Diphosphosulfoquinovose  
(UDP-SQ)**

Filing Date:

Serial No.:

WHEREAS, Michigan State University, a Michigan corporation, 238 Administration Building, East Lansing, Michigan 48824-1046, hereinafter referred to as "ASSIGNEE", is desirous of acquiring the entire right, title and interest in said invention and application and in any Letters Patent which may be granted on the same;

NOW THEREFORE, TO ALL WHOM IT MAY CONCERN: Be it known that, for and in consideration of the sum of One Dollar (\$1.00) lawful money paid to ASSIGNOR by ASSIGNEE and other good and valuable consideration, receipt of which is hereby acknowledged, ASSIGNOR has sold, assigned and transferred, and by these presents does sell, assign and transfer unto said ASSIGNEE, and ASSIGNEE's successors and assigns, all right, title and interest in and to said invention, said application for United States Letters Patent, and any Letters Patent which may hereafter be granted on the same in the United States and all countries throughout the world including any divisions, renewals, continuations in whole or in part, substitutions, conversions, reissues, prolongations or extensions thereof, said interest to be held and enjoyed by said ASSIGNEE as fully and exclusively as it would have been held and enjoyed by said ASSIGNOR had this assignment and transfer not been made, to the full end and term of any Letters Patent.

ASSIGNOR further agrees that ASSIGNOR will, without charge to said ASSIGNEE, but at ASSIGNEE's expense, cooperate with ASSIGNEE in the prosecution of said application and/or applications; execute, verify, acknowledge and deliver all such further papers, including applications for Letters Patent and for the reissue thereof, and instruments of assignment and transfer thereof; and will perform such other acts as ASSIGNEE lawfully may request, to obtain or maintain Letters Patent for said invention and improvement in any and all countries, and to vest title thereto in said ASSIGNEE, or ASSIGNEE's successors and assigns.

IN TESTIMONY WHEREOF, ASSIGNOR has hereunto signed ASSIGNOR's names to this assignment on the date indicated below.

\_\_\_\_\_  
Christoph Benning

\_\_\_\_\_  
Sherrie Lea Sanda

\_\_\_\_\_  
Bin Yu

STATE OF \_\_\_\_\_)

COUNTY OF \_\_\_\_\_)

ss.

On this \_\_\_\_\_ day of \_\_\_\_\_, in the year of \_\_\_\_\_, before me, the undersigned notary public, personally appeared the above-named ASSIGNOR, known to me (or proved to me on the basis of satisfactory evidence) to be the person whose name is subscribed to the within instrument, and acknowledged that he/she executed the same.

\_\_\_\_\_  
NOTARY PUBLIC

SEAL



# SEQUENCE LISTING

<110> Benning, Christoph

Sanda, Sherrie

Yu, Bin

<120> Compositions and Methods for the Synthesis and Subsequent Modification of Uridine-5-Diphosphosulfoquinovose (UDP-SQ)

<130> MSU-04769

<160> 21

<170> PatentIn version 3.0

<210> 1

<211> 1134

<212> DNA

<213> Cyanobacterium Synechococcus

<220>

<221> CDS

<222> (1) .. (1134)

<400> 1	
atg cgc atc gct ctc ttt acc gag acg ttc ctc ccc aaa gtg gat ggc	48
Met Arg Ile Ala Leu Phe Thr Glu Thr Phe Leu Pro Lys Val Asp Gly	
1 5 10 15	
atc gtc acg cgg ctt cgg cac acg gtc gat cac ctg cag cgt ctt ggc	96
Ile Val Thr Arg Leu Arg His Thr Val Asp His Leu Gln Arg Leu Gly	
20 25 30	
cac acc gtc atg gtt ttt tgc ccc gac ggc ggg ctc cgc gag cac aag	144
His Thr Val Met Val Phe Cys Pro Asp Gly Gly Leu Arg Glu His Lys	
35 40 45	
ggg gct cga gtc tat ggg gtt aaa ggc ttt ccg cta ccg ctc tat ccc	192
Gly Ala Arg Val Tyr Gly Val Lys Gly Phe Pro Leu Pro Leu Tyr Pro	
50 55 60	
gag ctg aag cta gct ttt ccg ttg ccg aaa gtg gga aaa gcc ttg gag	240
Glu Leu Lys Leu Ala Phe Pro Leu Pro Lys Val Gly Lys Ala Leu Glu	
65 70 75 80	
cgg ttc cgg ccc gac ctg atc cac gtg gtc aat ccg gct gtg ttg ggg	288
Arg Phe Arg Pro Asp Leu Ile His Val Val Asn Pro Ala Val Leu Gly	
85 90 95	
ttg ggc ggc atc tac tat gcc aag gcg cta aat gtg cca ctc gtg gcg	336
Leu Gly Gly Ile Tyr Tyr Ala Lys Ala Leu Asn Val Pro Leu Val Ala	
100 105 110	

tcc Ser	tat Tyr	cac His 115	acc Thr	cat His	ttg Leu	ccg Pro	aaa Lys 120	tac Tyr	ctt Leu	gag Glu	cat His	tac Tyr 125	ggg Gly	ctg Leu	ggg Gly	384
gtc Val	ttg Leu 130	gag Glu	ggg Gly	gtg Val	ctc Leu	tgg Trp 135	gaa Glu	ttg Leu	ctg Leu	aag Lys	ctg Leu 140	gcg Ala	cat His	aac Asn	caa Gln	432
gca Ala 145	gcg Ala	atc Ile	aac Asn	ctc Leu	tgt Cys 150	act Thr	tca Ser	acc Thr	gcg Ala	atg Met 155	gtg Val	cag Gln	gag Glu	ctg Leu	aca Thr 160	480
gat Asp	cac His	ggc Gly	att Ile	gag Glu 165	cac His	tgt Cys	tgc Cys	ctc Leu	tgg Trp 170	cag Gln	cga Arg	gga Gly	gtg Val	gat Asp 175	acc Thr	528
gag Glu	acc Thr	ttt Phe	cgg Arg 180	cca Pro	gac Asp	ttg Leu	gct Ala 185	act Thr	gct Ala	gcg Ala	atg Met	cgc Arg	gat Asp 190	cgc Arg	ctc Leu	576
agt Ser	ggc Gly	ggg Gly 195	aag Lys	ccc Pro	act Thr	gcg Ala	ccc Pro 200	ttg Leu	ttg Leu	ctc Leu	tac Tyr	gtc Val 205	gga Gly	cgc Arg	ctc Leu	624
tca Ser	gcc Ala 210	gag Glu	aag Lys	caa Gln	atc Ile	gat Asp 215	cgc Arg	ctg Leu	cga Arg	ccc Pro	att Ile 220	ttg Leu	gat Asp	gcc Ala	aat Asn	672
cct Pro 225	gag Glu	gct Ala	tgc Cys	ttg Leu	gcc Ala 230	ttg Leu	gtc Val	ggc Gly	gat Asp	ggc Gly 235	ccg Pro	cat His	cgg Arg	gcc Ala	gaa Glu 240	720
cta Leu	gag Glu	caa Gln	ttg Leu	ttt Phe 245	gct Ala	ggc Gly	acc Thr	cag Gln	acg Thr 250	cag Gln	ttc Phe	att Ile	ggc Gly	tat Tyr 255	ctg Leu	768
cat His	ggg Gly	gaa Glu	cag Gln 260	cta Leu	ggg Gly	gcg Ala	gcc Ala	tac Tyr 265	gct Ala	tct Ser	gct Ala	gac Asp 270	gct Ala	ttt Phe	gtc Val	816
ttt Phe	ccc Pro	tcc Ser 275	cgg Arg	acc Thr	gaa Glu	acc Thr	ctc Leu 280	ggg Gly	cta Leu	gtc Val	ttg Leu	ctg Leu 285	gaa Glu	gcc Ala	atg Met	864
gca Ala 290	gcg Ala	ggg Gly	tgt Cys	ccg Pro	gtc Val	gtg Val 295	gcg Ala	gcc Ala	aat Asn	tcc Ser	ggg Gly 300	ggc Gly	att Ile	ccc Pro	gat Asp	912
att Ile 305	gtc Val	agc Ser	gac Asp	ggc Gly	att Ile 310	aat Asn	ggg Gly	ttc Phe	ctg Leu	ttc Phe 315	gat Asp	cct Pro	gag Glu	gat Asp	gaa Glu 320	960
caa Gln	ggg Gly	gcg Ala	atc Ile	gct Ala 325	gcg Ala	att Ile	cag Gln	cgc Arg	ttg Leu 330	ttg Leu	gct Ala	aac Asn	cct Pro	gca Ala 335	gag Glu	1008
cgc Arg	gag Glu	att Ile	cta Leu 340	cgc Arg	caa Gln	gcg Ala	gct Ala 345	cgt Arg	caa Gln	gaa Glu	gcc Ala	gaa Glu	cgc Arg 350	tgg Trp	agc Ser	1056



Glu Thr Phe Arg Pro Asp Leu Ala Thr Ala Ala Met Arg Asp Arg Leu  
 180 185 190  
 Ser Gly Gly Lys Pro Thr Ala Pro Leu Leu Leu Tyr Val Gly Arg Leu  
 195 200 205  
 Ser Ala Glu Lys Gln Ile Asp Arg Leu Arg Pro Ile Leu Asp Ala Asn  
 210 215 220  
 Pro Glu Ala Cys Leu Ala Leu Val Gly Asp Gly Pro His Arg Ala Glu  
 225 230 235 240  
 Leu Glu Gln Leu Phe Ala Gly Thr Gln Thr Gln Phe Ile Gly Tyr Leu  
 245 250 255  
 His Gly Glu Gln Leu Gly Ala Ala Tyr Ala Ser Ala Asp Ala Phe Val  
 260 265 270  
 Phe Pro Ser Arg Thr Glu Thr Leu Gly Leu Val Leu Leu Glu Ala Met  
 275 280 285  
 Ala Ala Gly Cys Pro Val Val Ala Ala Asn Ser Gly Gly Ile Pro Asp  
 290 295 300  
 Ile Val Ser Asp Gly Ile Asn Gly Phe Leu Phe Asp Pro Glu Asp Glu  
 305 310 315 320  
 Gln Gly Ala Ile Ala Ala Ile Gln Arg Leu Leu Ala Asn Pro Ala Glu  
 325 330 335  
 Arg Glu Ile Leu Arg Gln Ala Ala Arg Gln Glu Ala Glu Arg Trp Ser  
 340 345 350  
 Trp Asn Ala Ala Thr Arg Gln Leu Gln Asp Tyr Tyr Cys Glu Val Leu  
 355 360 365  
 Ala Asp Gly Cys Leu Pro Leu Ala Ala  
 370 375

<210> 3

<211> 2979

<212> DNA

<213> Arabidopsis thaliana

<400> 3  
ctacacgtta ccttccggta ctggaaacag tcgtttaatc aaccaattga ttgggtcccaa 60  
aacatgaact ttctttttcc tccagaacca aatcgctgca ctgtactgtt cattgcgtat 120  
ctttgtcgtt gctgctctcc aatcatatct ctcggtctct tctcttgccg cttttccaat 180  
gatctctctt gtttcacggc cgtgcagtaa agttctcagt tttgtcacgc aatcttcaac 240  
atctccaggg ttgaacaaaa atccgggttt tccctcctga tgaaaacatc agaatcagaa 300  
aaccacaagc tcaatatagg ttgaccata agaacaatca atgcaagatc attttgtgta 360  
ccagtctatg attgaataaa gtttcagttc gggttacagc cgcttataag aaaattggca 420  
gaaattgttt tttcaaccat ttcgggttcg ttgatatgct catcaatatg gtttggcagt 480  
taattgtaat tcagataatt cactgacctg atcttcaggg attatatcag ggattccacc 540  
ggcacggggc gcgacgacag gaagtcctga agacattgct tcaagaacca caaggccaag 600  
tgtctccgac tctgatggca tcacaaacac atctccactt gcgtaagctt gtgagagtcc 660  
atcgccctgt aacgttccag tgaaaaccgc tggcattccg gtaaacaaact tctcaagatc 720  
ctctctttta gaaaacgaaa cagataaaca aaattacaat gttgttgact agaaatcttc 780  
agataacaat atggccaatc ttttaacaaa ctagtacttg tatgggtccat ctccaatgaa 840  
agcaatccga gtttcaggta atttgtccat tacactgcac acaaatttct caatatcaaa 900  
attcgataca ccacttaaaa gaagtgaagc cagtttatac aaaattctaa cctctttaaa 960  
agctccaaac tcttttctac gccaatgcga cctacatgaa tcactagtgg cttttctggt 1020  
tcgccattac tgtaattac aaaatattaa acatcaagat tagcgtggaa agtatcattg 1080  
tttttaaatgc atataaaaga aacgtatatt ctattcttgc ctcagtctta tacgcatttc 1140  
ttgagaacgg aaacggggat tgaagctttc tgaatcgaca cccttattcc aaagtcgaag 1200  
ttgattagct tcaatatatg aaaaaaagaa gaagaaaatg taagttttga acaatcatag 1260  
agcttgatca aaaatgtaat gtatgatctt tcttcacctg cagttgcacc agctgctata 1320  
agatcttttc caatggcagc agaaggaact aatgtaagat cagccgctct gtgaaggaac 1380  
cctgataaaa gcatattcaa gtttagtttc atattataca tcacaataaa ccagaaaaag 1440  
aagaaggaaa ttttgacatt tgaaaagcgg gttttacata cttattatag accacattgg 1500  
ttttaccaac caactaaaag tgtatcttgg gatgtatctg cacatcaaca caagcttaat 1560  
cttagacaaa aattttttta taacaacatt gtgaaatgag gcagaaaaag gtacttacac 1620  
aggacgtgt gtgtggtaag acattactat tggtagagat agcatttttg ctattgccag 1680  
agcaccaaag acctaaaaat ttagtcagg gaaaaaagag agtcaagttc tggattctct 1740  
ccagttcact ggtcttgcta atgttttagt aatgtgaatt cttgagggat ttaccataac 1800  
tccgggagat gaagcgtgta taatgtcagg cttaaaccgt gcaatttcag agatgattct 1860

tggactaagc gcaagcgaga gtggaacctt ttggaataa ggacaaggga agctacagaa 1920  
gagaagaaga aattagcgat attaccaaatt agagaacatc cagttagtaa actaaatggt 1980  
gctaccttct tgatccaatg actctggctc cataaaactc ttcaggaaca ctttcatgtg 2040  
tcgtcacgac tataacctat gaagcaaaaa agttattaaa aaaaaaaaaa aaggaaacagt 2100  
taacacttgt caagtaattc taattctgga aacagttact tatgagctga ctgaaaagat 2160  
acttaagttg aagaatgaga tagtaaaaga agaaacctcg tctcccattt cacggaggta 2220  
tctaataaaa ttctggaatc tgtttttgta gccggatata tagctgcaac aaaatcaaag 2280  
agagaatcac ttccaataat aacatgacat ataataaaag ctttttgtca atggatcggg 2340  
gattccgaga atcttgggat attcacaact aaaatctgac aactttgact caaacaatc 2400  
ctgaatgtaa ttggttttaa cgatctacta tataatttgc taaattgggtg gtgtagcaaa 2460  
ttcatacatt agcgagtatc tcttcataaa ataaatgtaa cgatcaaact gaaagaaaaa 2520  
aacattacag gaaaaagtca accaaggaaa aaaatgagta gaatctgttt tcacagagac 2580  
atctcgtcga acacaaaaca agcaaaaaaa gaacactgtg aagaagactt acgcaaaggg 2640  
agaaggctca acaaagagag caattctcct aggcttagag agcgactcag gatcgagaag 2700  
cggcgcacat atctccgatt cgtcatcttc tctgacttga gtaatagtca tatcggttga 2760  
cccagaaaaca gcttcctttg taatcactcc acagaagcgg agcttgcttt tcttgctaatt 2820  
gggtaatctc cggtgaccaa acgaaagagg ggaatgaaga acaaaagaag aagacctggg 2880  
aggagaacaa gaggttgacg aggaagaaga acaagtgtta gtcgtgctag gaagcaaattg 2940  
aggaggtata gagagattta tagaagaaag agtcgtcat 2979

<210> 4

<211> 2537

<212> DNA

<213> Arabidopsis thaliana

<400> 4

tcataatttg aaaaagcact tcttcttctt cttcttcttc ttcttcttct tcttcttccc 60  
atcaaaaagt gctagtggca agaccgcgca tattccagta acaaagttaa aattcaaaac 120  
tatggaacac aatttgctcc atgttttggt gcctgaatta aacgaccata tcaattgggtt 180  
cggccacttc atctcggata cgcacaagcg gttgtcaagg ttgtaaaaga cgattttgta 240  
aggatctaca ttggcaaagg gagctttaca gacgacttga aaagtctcgg tgtgaagatt 300  
gaaaggtaaa actttgggtt ctttgacttc ggtgaaccaa taaagcgacc catctagata 360  
cacagggtcg gggtaagcag caatccgata aggagcagcg agagtgcacat acctccaagc 420

attggtacta	aagtcgaaaa	cttcgcatgt	agtagcgttt	tttaggccta	tttctgcaga	480
gttgataaac	caaacgggct	tgtatgtgcc	cctgaatttg	tctttaccga	atccaagcat	540
aaagaatttg	cgtttaagct	tgtagtaacc	atctcgtaag	tcgatcatga	gtttttgata	600
atcgcaaaga	ggaagaggcc	gataccatct	agtggtcggg	ttgaccacat	aaccggattt	660
gtcgtgattg	taaagacaaa	cgagaccgtc	acaactacta	tgtgaaacta	agtacagtac	720
gttatccttc	tcccaaggag	tagggatctt	gaacgttgat	gatgaaccga	actgcaatgt	780
tcttaaagat	tctatggtcg	ggttttataac	atggtgagga	aatacagaca	ccatcagaac	840
atctggatct	cctgattgct	gacgatgttt	caactaaaat	tgtatacatt	tttattttat	900
taaaaagata	aatatcattt	tgttggaatt	ggaaaacaaa	atctaaaaga	taactttaga	960
tatgtaaggg	tcgcatatgt	atttgattac	attgctatga	gtatatgact	cacttttcca	1020
ataatgaaaa	aaataaagat	gttgagtctt	actaattaag	ggtaaataca	aaattttctga	1080
tcattaatac	aaagaaaaag	cctctcataa	gccaaagcca	ttttactcgc	cgtaaacatc	1140
tccgcagcat	actctttaca	cttccttcct	ctctccgcca	atctctccgc	tccttcagcc	1200
acagcaacct	ccataaccgc	cgtcaaagct	tccacatttg	gcgcaaacat	aaacccaaac	1260
tcatcattca	ccactatagt	cctctttatg	cttgcgtagc	tagatgccat	cactggttta	1320
ccactcaaca	tcgcttccat	taacgttaaa	tcaagacctt	gtggctctaag	cgttggaattc	1380
acgaacaaat	cgatcccgtt	gtagaaaccc	ttgagctcat	ttgggtttag	agatcccaaa	1440
atggaaaactt	tttcccctaa	ttccttgtaa	cgttgctccc	atggccctga	tccagctact	1500
acgaggtaaa	catttgaata	cgtttggatt	atcttcgcga	aagcttcgaa	gagcaatgga	1560
tgtcctttgt	ctttgactaa	tctcccagca	gctcctaaaa	caatcgctga	tgagttttct	1620
ggtaacccta	atcttgacct	aaacagagta	cgtagcttct	tgtctgatgt	gaatccgttc	1680
tcgtcgactc	cattgaggat	cacatgaacc	cttttctcag	ggatttggtg	aacgtctctt	1740
agcatttctc	cgcagctatc	gctgatagcg	atgtggtgag	cgtagttgtg	gaagaatctg	1800
atttcgtcca	gtatcttggg	aagcacggca	ccgtatagac	ttgcattgaa	gccttgatgat	1860
cttggttcgt	ctgggtttacg	gatcaggtct	tggtaaatac	ttgattgtaa	gctctctaac	1920
gcaatgccgt	gccaggatac	agcgaggttt	ggaacctccc	ggcgatccga	gtgaggtaaa	1980
gcaacacttt	cagagtgaac	cgcacgaaa	ggttctttct	tgttttcttc	ttggtaaagc	2040
tcccatgcct	tggtgtaccg	ccattttccg	ggctccgcgt	cgccgtgaga	atggattata	2100
gggtaaatga	tttggtcgga	aacagggggg	atcttggttg	tttcagggga	ttggtctaag	2160
ggagaggtga	aaacgtggac	acggtgtcca	cggcgagcta	aggcgggtga	tagagtgaag	2220
gcgtggcggt	ccatgccgcc	aggattggga	cccgtggggc	atttcgggga	gaaaacagcg	2280

agtttttagtg ttttgggagg agggttgggt aaggagaaat caaggcgggt ccaggcaaat 2340  
 tgggcgggtt ggaggtcgcc agaccacggt ggctggtttg tgtcagagga ggaggaggag 2400  
 actgcagcgg tggaggagga gcaagtggag gtgcggagga gaaagagagc cggtatggta 2460  
 aagaggacgg tgaagaagag gaaagtacaa aggctaaaat gagaattagg tttcttgagt 2520  
 ttggtttggtg aagccat 2537

<210> 5

<211> 2369

<212> DNA

<213> Arabidopsis thaliana

<400> 5

ttaaggtctc atgcatttga ccagaacatc gacgaatcgc ttgtacatat gttgcttcat 60  
 gtacatttttc tcaaccattt tgcgtccttc actccccaat cgtagcctct cgtctggatt 120  
 cctaagtaga tacaagagat tatgagctaa ttctttgtta ccggatctcc ccattgagtg 180  
 aagtagacca gtcatgttgt gttgaacat ctctttgggt cctcctgcat ctgttccac 240  
 cactgcaagt ccataagcca ttgcttcgat tgtcactcta ccaaagtgtt caccaactcc 300  
 ctgtaaaaaa tcaaggttca aaataaataa accgaacaaa atcgaaatgg tcaaaccgaa 360  
 atcaaacct acattttgaa caaaaacaaa actgaaatgt tttatccaaa accaaactga 420  
 aatcaatcac aaattgggtt atttgattca gtaattcaag ttcctataaa actgataaaa 480  
 ctaaaccaaa acctaactgt aaactacttt ctagtggaaa actgcatatg catcaaataa 540  
 tgtttttgag gtgtgaggag gattacctgg gagtttgtaa cgtagacatc tgctgcagag 600  
 tataatgaag caacacgggt tgttcgagga gtccacatta ccgacttaga taagtttccg 660  
 ctgtttgaca agaagcttaa catctcttta acgtatccaa ctttggttgct ctttgaaccc 720  
 acggatccta aaagaacttt aagttcttgc ttctctcttc ttaagccatt gtcaagtgtg 780  
 agagaaacac ttttcatctg tcgtgatgaa cctcttaaac gatgtttgct ggagagacta 840  
 accttttctt tcctaattgat ccccttgatg ttcccttgag attcttgtcc tctctcagaa 900  
 agagccaagg caatagattc aaggagaaga agttgtccct ttggtgggtt tatgctgcta 960  
 agagacatca caagcatatc tgaatctggt attcctaact ctgttctcac tgattcgcgt 1020  
 aatatttgtc tottcaacct cattttctct ggtgaaagtg ttggtgtggt gagtgaagaa 1080  
 ggaatccccg ctacaaaagc taactcatca ttaacagata gtggaacaat cactggttgt 1140  
 gatctaagct ttatatgctc ctctctgcac catgttaacc attgtctgct ctgtgattca 1200  
 gataagaaaa tcagcatttt cactcggatc agaactgggt tcgctcgatc aaagtattct 1260



```

cgtcgattct ccattatcca ccaagctatt tgacttccac cagctggatg atgattcatg 1320
tattgatctg cgagaaaagg aaaaaaaaaa agtaattaag atatgctttc ctctgattgt 1380
aaattagtag tctaaaaact gcataatgaa ttcttaccta tccatgaggt acacactgct 1440
gatcctgcga tgatcaaadc tgctttcatg gcagtcttga agctgagttc tcctttgtct 1500
tcaacaactt tgatccttct cctactgagc tcttgcacat atccacctct cctgctaaga 1560
actactgcag agactggtgc accacagctc aaaagctctg aagccagctc catcatagaa 1620
attggagcac cagtcattga tagctcgtgg aaaagcagga cgaatctcct tgaccaaaaca 1680
aggcggttaa aatctgattt cctgtcgcaa gtcccagatc ttctatgcgg gctccattca 1740
agaactttat cctctagtga tccaaaggga ccaagaagct taccataagt agcatttgtc 1800
aatggaagtt gtggatcttg ctcatcatcc aaatccttag tctcaagtac ctctttaatc 1860
accttctgct taacacggat cttactacga gaagtccgga cagtttttct cgtcttctgc 1920
ttggaactca agctccgctg agaaacccca tcatctttct tgatcagact aacatccgct 1980
ctcttattcg aaccagcatc atctttacca gtaatatata ccaaagcctc agaattctca 2040
ttagcaacaa catccagtcc tttaatcttc tccatatata actcatcctt tctcggctctg 2100
cctccaaacc gtaaaaaactc aactttgctt tcattatcat gtgccacact agactgaaca 2160
taaaatccaa gatacgtcca aagcgtaatc aaaagcagcc aataaaccaa ccggctacta 2220
cgaaaccact gaaacgctcc tctccacca tgacctctac ggggagtcct accagaatac 2280
actctaggtg tacccttgg agtagacctc cctgacagtg aagacttaac acttgtctgt 2340
ctcagcgggc ataaccgaat ctctccat 2369

```

<210> 6

<211> 1799

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (170)..(1603)

```

<400> 6
gtcgaccac gcgtccgctc atctctcatc gttccgggag aagagaagag agacccatcc 60
ctcacttcaa agttcaaagt ctccaaggat cttctccaac tctctctaaa caagattcca 120
aattttcaaa ggtgaatttg ttgatagaa tcaagaacaa acctttaaa atg gcg cat 178
Met Ala His
1
cta ctt tca gct tca tgc cct tca gtt atc tca ctt agc agc agc agc 226

```

Leu	Leu	Ser	Ala	Ser	Cys	Pro	Ser	Val	Ile	Ser	Leu	Ser	Ser	Ser	Ser	
5						10					15					
agc	aag	aat	tca	gtt	aag	ccg	ttt	gtt	tca	ggg	cag	acc	ttc	ttc	aat	274
Ser	Lys	Asn	Ser	Val	Lys	Pro	Phe	Val	Ser	Gly	Gln	Thr	Phe	Phe	Asn	
20					25					30					35	
gct	cag	ctt	ctt	tca	aga	tct	tct	ctc	aaa	gga	ctt	ctc	ttc	caa	gag	322
Ala	Gln	Leu	Leu		Arg	Ser	Ser	Leu	Lys	Gly	Leu	Leu	Phe	Gln	Glu	
				40					45					50		
aag	aaa	ccg	aga	aaa	agc	tgc	gtt	ttc	aga	gca	act	gct	gta	cct	ata	370
Lys	Lys	Pro	Arg	Lys	Ser	Cys	Val	Phe	Arg	Ala	Thr	Ala	Val	Pro	Ile	
			55					60					65			
acc	caa	caa	gca	cca	ccc	gaa	aca	tct	acc	aat	aac	tca	tcc	tct	aaa	418
Thr	Gln	Gln	Ala	Pro	Pro	Glu	Thr	Ser	Thr	Asn	Asn	Ser	Ser	Ser	Lys	
		70					75					80				
cca	aag	cgt	gtt	atg	gtc	att	ggg	gga	gat	ggg	tat	tgc	ggg	tgg	gct	466
Pro	Lys	Arg	Val	Met	Val	Ile	Gly	Gly	Asp	Gly	Tyr	Cys	Gly	Trp	Ala	
	85					90					95					
act	gct	ctc	cac	ttg	tcc	aag	aag	aat	tac	gaa	gtt	tgc	att	gtt	gac	514
Thr	Ala	Leu	His	Leu	Ser	Lys	Lys	Asn	Tyr	Glu	Val	Cys	Ile	Val	Asp	
100					105					110					115	
aac	ctt	gta	aga	cgt	ctt	ttc	gac	cac	cag	ctt	gga	ctt	gag	tca	ttg	562
Asn	Leu	Val	Arg	Arg	Leu	Phe	Asp	His	Gln	Leu	Gly	Leu	Glu	Ser	Leu	
				120					125					130		
act	cct	att	gcc	tcc	att	cat	gac	cga	atc	agc	cga	tgg	aag	gct	ttg	610
Thr	Pro	Ile	Ala	Ser	Ile	His	Asp	Arg	Ile	Ser	Arg	Trp	Lys	Ala	Leu	
			135					140					145			
aca	ggg	aaa	tca	att	gag	ttg	tac	gtt	ggg	gat	atc	tgt	gat	ttc	gaa	658
Thr	Gly	Lys	Ser	Ile	Glu	Leu	Tyr	Val	Gly	Asp	Ile	Cys	Asp	Phe	Glu	
		150					155					160				
ttc	tta	gct	gag	tct	ttc	aag	tct	ttt	gag	ccg	gat	tca	gtt	gtc	cac	706
Phe	Leu	Ala	Glu	Ser	Phe	Lys	Ser	Phe	Glu	Pro	Asp	Ser	Val	Val	His	
	165					170					175					
ttt	ggg	gaa	cag	aga	tcc	gct	cct	tac	tcg	atg	att	gac	cgg	tcc	aga	754
Phe	Gly	Glu	Gln	Arg	Ser	Ala	Pro	Tyr	Ser	Met	Ile	Asp	Arg	Ser	Arg	
180					185					190					195	
gca	gtt	tat	aca	cag	cac	aac	aat	gtg	att	ggg	act	ctc	aac	gtt	ctc	802
Ala	Val	Tyr	Thr	Gln	His	Asn	Asn	Val	Ile	Gly	Thr	Leu	Asn	Val	Leu	
				200					205					210		
ttt	gct	ata	aaa	gag	ttt	gga	gag	gag	tgt	cat	ctt	gta	aaa	ctt	ggg	850
Phe	Ala	Ile	Lys	Glu	Phe	Gly	Glu	Glu	Cys	His	Leu	Val	Lys	Leu	Gly	
			215					220					225			
acg	atg	ggg	gag	tat	gga	act	cca	aat	att	gac	atc	gag	gaa	ggg	tat	898
Thr	Met	Gly	Glu	Tyr	Gly	Thr	Pro	Asn	Ile	Asp	Ile	Glu	Glu	Gly	Tyr	
		230					235					240				
ata	acc	ata	acc	cac	aac	ggg	aga	act	gac	act	ttg	cca	tac	ccc	aag	946
Ile	Thr	Ile	Thr	His	Asn	Gly	Arg	Thr	Asp	Thr	Leu	Pro	Tyr	Pro	Lys	
		245				250					255					



<210> 7  
 <211> 477  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 7

Met Ala His Leu Leu Ser Ala Ser Cys Pro Ser Val Ile Ser Leu Ser  
 1 5 10 15

Ser Ser Ser Ser Lys Asn Ser Val Lys Pro Phe Val Ser Gly Gln Thr  
 20 25 30

Phe Phe Asn Ala Gln Leu Leu Ser Arg Ser Ser Leu Lys Gly Leu Leu  
 35 40 45

Phe Gln Glu Lys Lys Pro Arg Lys Ser Cys Val Phe Arg Ala Thr Ala  
 50 55 60

Val Pro Ile Thr Gln Gln Ala Pro Pro Glu Thr Ser Thr Asn Asn Ser  
 65 70 75 80

Ser Ser Lys Pro Lys Arg Val Met Val Ile Gly Gly Asp Gly Tyr Cys  
 85 90 95

Gly Trp Ala Thr Ala Leu His Leu Ser Lys Lys Asn Tyr Glu Val Cys  
 100 105 110

Ile Val Asp Asn Leu Val Arg Arg Leu Phe Asp His Gln Leu Gly Leu  
 115 120 125

Glu Ser Leu Thr Pro Ile Ala Ser Ile His Asp Arg Ile Ser Arg Trp  
 130 135 140

Lys Ala Leu Thr Gly Lys Ser Ile Glu Leu Tyr Val Gly Asp Ile Cys  
 145 150 155 160

Asp Phe Glu Phe Leu Ala Glu Ser Phe Lys Ser Phe Glu Pro Asp Ser  
 165 170 175

Val Val His Phe Gly Glu Gln Arg Ser Ala Pro Tyr Ser Met Ile Asp  
 180 185 190

Arg Ser Arg Ala Val Tyr Thr Gln His Asn Asn Val Ile Gly Thr Leu  
 195 200 205

00705030.1000

Asn Val Leu Phe Ala Ile Lys Glu Phe Gly Glu Glu Cys His Leu Val  
 210 215 220  
 Lys Leu Gly Thr Met Gly Glu Tyr Gly Thr Pro Asn Ile Asp Ile Glu  
 225 230 235 240  
 Glu Gly Tyr Ile Thr Ile Thr His Asn Gly Arg Thr Asp Thr Leu Pro  
 245 250 255  
 Tyr Pro Lys Gln Ala Ser Ser Phe Tyr His Leu Ser Lys Val His Asp  
 260 265 270  
 Ser His Asn Ile Ala Phe Thr Cys Lys Ala Trp Gly Ile Arg Ala Thr  
 275 280 285  
 Asp Leu Asn Gln Gly Val Val Tyr Gly Val Lys Thr Asp Glu Thr Glu  
 290 295 300  
 Met His Glu Glu Leu Arg Asn Arg Leu Asp Tyr Asp Ala Val Phe Gly  
 305 310 315 320  
 Thr Ala Leu Asn Arg Phe Cys Val Gln Ala Ala Val Gly His Pro Leu  
 325 330 335  
 Thr Val Tyr Gly Lys Gly Gly Gln Thr Arg Gly Tyr Leu Asp Ile Arg  
 340 345 350  
 Asp Thr Val Gln Cys Val Glu Ile Ala Ile Ala Asn Pro Ala Lys Ala  
 355 360 365  
 Gly Glu Phe Arg Val Phe Asn Gln Phe Thr Glu Gln Phe Ser Val Asn  
 370 375 380  
 Glu Leu Ala Ser Leu Val Thr Lys Ala Gly Ser Lys Leu Gly Leu Asp  
 385 390 395 400  
 Val Lys Lys Met Thr Val Pro Asn Pro Arg Val Glu Ala Glu Glu His  
 405 410 415  
 Tyr Tyr Asn Ala Lys His Thr Lys Leu Met Glu Leu Gly Leu Glu Pro  
 420 425 430  
 His Tyr Leu Ser Asp Ser Leu Leu Asp Ser Leu Leu Asn Phe Ala Val  
 435 440 445  
 Gln Phe Lys Asp Arg Val Asp Thr Lys Gln Ile Met Pro Ser Val Ser

460

<213> Cyanobacterium Synechococcus

<210> 9

<211> 402

<212> PRT

<213> Cyanobacterium Synechococcus

<400> 9

Met Lys Ile Leu Val Leu Gly Gly Asp Gly Phe Cys Gly Trp Pro Cys  
1 5 10 15  
Ala Leu Asn Leu Ala Ala Ala Gly His Ala Val Thr Ile Val Asp Asn  
20 25 30  
Leu Val Arg Arg Lys Thr Asp Val Glu Leu Gly Val Gln Ser Leu Thr  
35 40 45  
Pro Ile Ala Thr Ile Glu Arg Arg Leu Lys Ala Trp Gln Glu Thr Gly  
50 55 60  
Gly Gln Pro Ile Ser Phe Val Asn Leu Asp Leu Ala Ala Asp Tyr Asp  
65 70 75 80  
Arg Leu Cys Ala Leu Leu Leu Glu Thr Gln Pro Asp Ala Ile Val His  
85 90 95  
Phe Ala Glu Gln Arg Ala Ala Pro Tyr Ser Met Lys Ser Ala Trp His  
100 105 110  
Lys Arg Phe Thr Val Asn Asn Asn Val Asn Ala Thr His Asn Leu Leu  
115 120 125  
Cys Ala Cys Val Asp Val Gly Leu Lys Ser His Ile Val His Leu Gly  
130 135 140  
Thr Met Gly Val Tyr Gly Tyr Gly Ser His Arg Gly Ala Thr Ile Pro  
145 150 155 160  
Glu Gly Tyr Leu Glu Val Glu Val Val Gln Arg Asp Gly Gln Arg Phe  
165 170 175  
Glu Glu Lys Ile Leu His Pro Val Asp Pro Gly Ser Val Tyr His Met  
180 185 190  
Thr Lys Thr Leu Asp Gln Leu Leu Phe Tyr Tyr Tyr Asn Lys Asn Asp  
195 200 205  
Asn Ile Gln Val Thr Asp Leu His Gln Gly Ile Val Trp Gly Thr Asn  
210 215 220  
Thr Asp His Cys Asn Leu His Pro Asp Leu Thr Asn Arg Phe Asp Tyr  
225 230 235 240  
Asp Gly Asp Tyr Gly Thr Val Leu Asn Arg Phe Leu Met Gln Ala Ala  
245 250 255  
Ile Gly Tyr Pro Leu Thr Val His Gly Val Gly Gly Gln Thr Arg Ala  
260 265 270  
Phe Ile His Ile Arg Asp Ser Val Arg Cys Val Gln Leu Ala Ile Glu  
275 280 285

Asn Pro Pro Ala Ala Asn Glu Lys Val Arg Ile Phe Asn Gln Met Thr  
 290 295 300  
 Glu Thr Tyr Gln Val Lys Asp Leu Ala Glu Lys Val Ala Ala Leu Thr  
 305 310 315 320  
 Gly Ala Glu Ile Ala Tyr Leu Pro Asn Pro Arg Lys Glu Ala Leu Glu  
 325 330 335  
 Asn Asp Leu Ile Val Asp Asn Arg Cys Leu Ile Asp Leu Gly Leu Asn  
 340 345 350  
 Pro Thr Thr Leu Asp Asn Gly Leu Met Ser Glu Val Val Glu Ile Ala  
 355 360 365  
 Gln Lys Phe Ala Asp Arg Cys Asp Arg Ala Lys Ile Pro Cys Val Ser  
 370 375 380  
 Ala Trp Thr Arg Asn Gln Ala Glu Ala Leu Ser Ala Pro Glu Thr Ala  
 385 390 395 400  
 Leu Arg

<210> 10

<211> 26

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc\_feature

<223> Synthetic

<400> 10

aaaggatccc gtgttatggc cattgg

26

<210> 11

<211> 26

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc\_feature

<223> Synthetic

<400> 11

gtcggatcct tatgtgggtca tggact

26

<210> 12

<211> 24





aaggatccct acacgttacc ttccggta	28
<210> 16	
<211> 28	
<212> DNA	
<213> Artificial/Unknown	
<220>	
<221> misc_feature	
<223> Synthetic	
<400> 16	
aaggatccat ggcttcacaa accaaact	28
<210> 17	
<211> 28	
<212> DNA	
<213> Artificial/Unknown	
<220>	
<221> misc_feature	
<223> Synthetic	
<400> 17	
gcggatcctc atattttgaa aaagcact	28
<210> 18	
<211> 27	
<212> DNA	
<213> Artificial/Unknown	
<220>	
<221> misc_feature	
<223> Synthetic	
<400> 18	
agggtaccat ggagggattc ggttatc	27
<210> 19	
<211> 28	
<212> DNA	
<213> Artificial/Unknown	
<220>	

<221> misc\_feature  
 <223> Synthetic  
 <400> 19  
 gcggtacctt aaggtctatg catttgac 28  
 <210> 20  
 <211> 26  
 <212> DNA  
 <213> Artificial/Unknown  
 <220>  
 <221> misc\_feature  
 <223> Synthetic  
 <400> 20  
 ctaggtacca aatggcgcat ctactt 26  
 <210> 21  
 <211> 26  
 <212> DNA  
 <213> Artificial/Unknown  
 <220>  
 <221> misc\_feature  
 <223> Synthetic  
 <400> 21  
 aaggatcctg cgctaaagtc gcactc 26

## CLAIMS

We claim:

1. A method, comprising:

a) providing:

- i) uridine-5'-diphosphoglucose;
- ii) a sulfur donor;
- iii) a first peptide capable of catalyzing the conversion of uridine-5'-diphosphoglucose to uridine-5'-diphosphosulfoquinovose; and
- iv) a second peptide capable of transferring sulfoquinovose from uridine-5'-diphosphosulfoquinovose onto diacylglycerol;

b) reacting said uridine-5'-diphosphoglucose with said first peptide and said sulfur donor under such conditions that uridine-5'-diphosphosulfoquinovose is generated; and

c) treating said uridine-5'-diphosphosulfoquinovose with said second peptide under conditions such that sulfoquinovose diacylglycerol is generated.

2. The method of Claim 1, wherein said first peptide is encoded by the nucleic acid sequence set forth in SEQ ID NO: 6.

3. The method of Claim 1, wherein said second peptide is encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 5, and portions thereof.

4. The method of Claim 1, wherein said sulfur donor is selected from the group consisting of sulfate, sulfide, thiosulfate, sulfoglutathione, adenosine 5'-phosphosulfate, and 3'-phosphoadenosine-5'-phosphosulfate.

5. The method of Claim 1, wherein said sulfur donor is sulfite.

6. A method, comprising:

a) providing:

i) uridine-5'-diphosphoglucose;

ii) a sulfur donor;

iii) a peptide capable of catalyzing the conversion of uridine-5'-diphosphoglucose to uridine-5'-diphosphosulfoquinovose;

iv) an acid catalyst;

v) a short-chain alcohol; and

vi) a long-chain alcohol;

b) reacting said uridine-5'-diphosphoglucose with said peptide and said sulfur donor under such conditions that uridine-5'-diphosphosulfoquinovose is generated;

c) reacting said uridine-5'-diphospho-sulfoquinovose with said short-chain alcohol and said acid catalyst under such conditions that a short-chain alkyl sulfoquinovoside is generated;

d) treating said short-chain alkyl sulfoquinovoside with said long-chain alcohol under such conditions that a long-chain alkyl sulfoquinovoside is generated.

7. The method of Claim 6, wherein said short-chain alcohol is selected from the group consisting of methanol, ethanol, propanol, pentanol, hexanol, heptanol, octanol, nonanol, and isomers thereof.

8. The method of Claim 6, wherein said short-chain alcohol is butanol.

9. The method of Claim 6, wherein said acid catalyst is selected from the group consisting of  $\text{H}_2\text{SO}_4$ ,  $\text{HCl}$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{BF}_3$ , ortho-toluenesulfonic acid,

meta-toluenesulfonic acid, alkylbenzenesulfonic acid, secondary alkyl-sulfonic acid, sulfonic resin, alkylsulfate, alkylbenzenesulfonate, alkyl-sulfonate, and sulfosuccinic acid.

10. The method of Claim 6, wherein said acid catalyst is para-  
5 toluenesulfonic acid.

11. The method of Claim 6, wherein said long-chain alcohol is selected from the group consisting of n-dodecyl alcohol, n-tetradecyl alcohol, n-hexadecyl alcohol, n-octadecyl alcohol, n-octyl alcohol, n-decyl alcohol, undecyl alcohol, and tridecyl alcohol.

12. The long-chain alkyl sulfoquinovoside prepared according to Claim 6.

13. A method, comprising:

a) providing:

i) uridine-5'-diphosphoglucose;

ii) a sulfur donor; and

iii) a peptide encoded by the nucleic acid sequence set forth  
in SEQ ID NO: 5; and

b) reacting said uridine-5'-diphosphoglucose with said peptide and  
said sulfur donor under such conditions that uridine-5'-  
diphosphosulfoquinovose is generated.

14. The method of Claim 13, wherein said sulfur donor is sulfite.

## ABSTRACT

The present invention is directed to compositions and methods related to the synthesis and modification of uridine-5'-diphospho-sulfoquinovose (UDP-SQ). In particular, the methods of the present invention comprise the utilization of recombinant enzymes from *Arabidopsis thaliana*, UDP-glucose, and a sulfur donor to synthesize UDP-SQ, and the subsequent modification of UDP-SQ to form compounds including, but not limited to, 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol (SQDG) and alkyl sulfoquinovoside. The compositions and methods of the invention provide a more simple, rapid means of synthesizing UDP-SQ, and the subsequent modification of UDP-SQ to compounds including, but not limited to, SQDG.